arXiv:math/0601425v1 [math.QA] 18 Jan 2006

UNITARY REPRESENTATIONS OF THE EXTENDED AFFINE LIE ALGEBRA $\widetilde{\mathfrak{gl}_3(\mathbb{C}_q)}$

Ziting Zeng

Abstract. We present modules for the extended affine Lie algebra $\widetilde{\mathfrak{gl}_3(\mathbb{C}_q)}$ by using the idea of free fields. A necessary and sufficient condition for the modules being unitary is given.

§0. Introduction.

Let q be a non-zero complex number. A quantum 2-torus associated to q (see [M]) is the unital associative $\mathbb{C}$-algebra $\mathbb{C}_q[s^{\pm 1}, t^{\pm 1}]$ (or, simply $\mathbb{C}_q$) with generators $s^{\pm 1}, t^{\pm 1}$ and relations

$$ss^{-1} = s^{-1}s = tt^{-1} = t^{-1}t = 1 \text{ and } ts = qst. \tag{0.1}$$

Define $\kappa, : \mathbb{C}_q \to \mathbb{C}$ to be a $\mathbb{C}$-linear function given by

$$\kappa(s^m t^n) = \delta_{(m,n),(0,0)} \tag{0.2}$$

Let d_s, d_t be the degree operators on $\mathbb{C}_q$ defined by

$$d_s(s^m t^n) = m s^m t^n, \ d_t(s^m t^n) = n s^m t^n \tag{0.3}$$

for $m, n \in \mathbb{Z}$.

Let $\mathfrak{gl}_3(\mathbb{C}_q)$ be the Lie algebra of 3 by 3 matrices whose entries are from $\mathbb{C}_q$. We form a natural central extension of $\mathfrak{gl}_3(\mathbb{C}_q)$ as follows.

$$\widehat{\mathfrak{gl}_3(\mathbb{C}_q)} = \mathfrak{gl}_3(\mathbb{C}_q) \oplus \mathbb{C}c_s \oplus \mathbb{C}c_t \tag{0.4}$$

with Lie bracket

(0.5)

$$\begin{aligned}&[E_{ij}(s^{m_1}t^{n_1}), E_{kl}(s^{m_2}t^{n_2})]\\ =&\delta_{jk}q^{n_1m_2}E_{il}(s^{m_1+m_2}t^{n_1+n_2}) - \delta_{il}q^{n_2m_1}E_{kj}(s^{m_1+m_2}t^{n_1+n_2})\\ &+ m_1q^{n_1m_2}\delta_{jk}\delta_{il}\delta_{m_1+m_2,0}\delta_{n_1+n_2,0}c_s + n_1q^{n_1m_2}\delta_{jk}\delta_{il}\delta_{m_1+m_2,0}\delta_{n_1+n_2,0}c_t\end{aligned}$$

Typeset by $\mathcal{AMS}$-TeX

for $m_1, m_2, n_1, n_2 \in \mathbb{Z}$, $1 \leq i, j, k, l \leq 3$, where E_{ij} is the matrix whose (i,j)-entry is 1 and 0 elsewhere, and c_s and c_t are central elements of $\widehat{\mathfrak{gl}_3(\mathbb{C}_q)}$.

The derivations d_s and d_t can be extended to derivations on $\mathfrak{gl}_3(\mathbb{C}_q)$. Now we can define the semi-direct product of the Lie algebra $\widehat{\mathfrak{gl}_3(\mathbb{C}_q)}$ and those derivations:

$$\widetilde{\mathfrak{gl}_3(\mathbb{C}_q)} = \widehat{\mathfrak{gl}_3(\mathbb{C}_q)} \oplus \mathbb{C}d_s \oplus \mathbb{C}d_t. \tag{0.6}$$

The Lie algebra $\widetilde{\mathfrak{gl}_3(\mathbb{C}_q)}$ is an extended affine Lie algebra of type A_2 with nullity 2. (See [AABGP] and [BGK] for definitions).

Extended affine Lie algebras are a higher dimensional generalization of affine Kac-Moody Lie algebras introduced by [H-KT] and systematically studied in [AABGP] and [BGK]. It turns out that any extended affine Lie algebra of type A is coordinated by a quantum torus (or a nonassociative torus for some small rank cases). Representations for extended affine Lie algebras coordinated by quantum tori and Lie algebras related to quantum tori have been studied in [JK2], [BS], [G1,2,3], [ER1,2], [EB],[GZ] [EZ], [LT1,2], [G-KK], [VV], [Mi], [ZZ], [BZ], [SZ], [L] and [BEG], and among others.

The Wakimoto's free fields construction provides a remarkable way to realize affine Kac-Moody Lie algebras (see [W2], [FF] and [EFK]). In [GZ], we used Wakimoto's idea to construct a class of representations for $\widetilde{\mathfrak{gl}_2(\mathbb{C}_q)}$ and found out the necessary and sufficient condition for the representations being unitary. In this paper, we will continue to construct representations for $\widetilde{\mathfrak{gl}_3(\mathbb{C}_q)}$. As witnessed in [FF], the realization for $\widetilde{\mathfrak{gl}_3(\mathbb{C}_q)}$ is much more subtle and complicated than the one for $\widetilde{\mathfrak{gl}_2(\mathbb{C}_q)}$. We then go on to construct a hermitian form and to determine when the form is positive definite (so the representations are unitary). Unlike [GZ] in which we defined the form on the monomial basis for the module (this idea goes back to [W1]), we define the form directly on the basis consisting of certain iterated module actions on a "highest weight vector" 1. This facilitates the verification of the defined form being a hermitian from.

Throughout this paper, we denote the field of complex numbers, real numbers and the ring of integers by $\mathbb{C}$, $\mathbb{R}$ *and* $\mathbb{Z}$ *respectively.*

§1. Module for $\widetilde{\mathfrak{gl}_3(\mathbb{C}_q)}$.

In this section, we use Wakimoto's idea [W1] to construct $\widetilde{\mathfrak{gl}_3(\mathbb{C}_q)}$-modules as was done in [GZ].

Let $\mathbb{K}_1 = \{(3m+1, 3n+1), m, n \in \mathbb{Z}\}$, and $\mathbb{K}_{-1} = \{(3m-1, 3n-1), m, n \in \mathbb{Z}\}$. If $\boldsymbol{A} = (3m+1, 3n+1) \in \mathbb{K}_1$, we always write $\boldsymbol{A}_1 = m, \boldsymbol{A}_2 = n$, and similarly, if $\boldsymbol{B} = (3m-1, 3n-1) \in \mathbb{K}_{-1}$, then $\boldsymbol{B}_1 = m, \boldsymbol{B}_2 = n$. Set

$$V = \mathbb{C}[x_{\boldsymbol{A}}, x_{\boldsymbol{B}} : \boldsymbol{A} \in \mathbb{K}_1, \boldsymbol{B} \in \mathbb{K}_{-1}] \tag{1.1}$$

be the (commutative) polynomial ring of infinitely many variables. The operators $x_{(m,n)}$ and $\frac{\partial}{\partial x_{(m,n)}}$ act on V as the usual multiplication and differentiation operators respectively.

Given a family of 2×2 lower triangular matrices

$$X_{m,n} = \begin{pmatrix} a_{(m,n)} & 0 \\ c_{(m,n)} & d_{(m,n)} \end{pmatrix} \in \ \mathrm{SL}_2(\mathbb{C})$$

for $(m,n) \in \mathbb{K}_1 \bigcup \mathbb{K}_{-1}$ (so $a_{(m,n)}d_{(m,n)} = 1$).

Set

$$P_{(m,n)} = a_{(m,n)} \frac{\partial}{\partial x_{(m,n)}} \tag{1.2}$$

$$Q_{(m,n)} = c_{(m,n)} \frac{\partial}{\partial x_{(m,n)}} + d_{(m,n)} x_{(m,n)} \tag{1.3}$$

for $(m,n) \in \mathbb{K}_1 \bigcup \mathbb{K}_{-1}$. Then for $\boldsymbol{A}, \boldsymbol{A'} \in \mathbb{K}_1, \boldsymbol{B}, \boldsymbol{B'} \in \mathbb{K}_{-1}$,

$$\begin{aligned} &[P_{\boldsymbol{A}}, P_{\boldsymbol{A'}}] = [Q_{\boldsymbol{A}}, Q_{\boldsymbol{A'}}] = [P_{\boldsymbol{B}}, P_{\boldsymbol{B'}}] = [Q_{\boldsymbol{B}}, Q_{\boldsymbol{B'}}] = 0 \\ &[P_{\boldsymbol{A}}, P_{\boldsymbol{B}}] = [P_{\boldsymbol{A}}, Q_{\boldsymbol{B}}] = [Q_{\boldsymbol{A}}, Q_{\boldsymbol{B}}] = [P_{\boldsymbol{B}}, Q_{\boldsymbol{A}}] = 0 \\ &[P_{\boldsymbol{A}}, Q_{\boldsymbol{A'}}] = \delta_{\boldsymbol{A},\boldsymbol{A'}}, \quad [P_{\boldsymbol{B}}, Q_{\boldsymbol{B'}}] = \delta_{\boldsymbol{B},\boldsymbol{B'}}. \end{aligned}$$

Fix a complex number μ, we define the following operators on V:

(1.4)

$$\begin{aligned} e^{(\mu)}_{21}(m_1,n_1) = &- q^{-m_1 n_1} \mu P_{-(3m_1-1,3n_1-1)} \\ &- \sum_{\boldsymbol{A},\boldsymbol{A'} \in \mathbb{K}_1} q^{n_1 \boldsymbol{A}'_1 + \boldsymbol{A}_2 m_1 + \boldsymbol{A}_2 \boldsymbol{A}'_1} Q_{\boldsymbol{A}+\boldsymbol{A'}+(3m_1-1,3n_1-1)} P_{\boldsymbol{A}} P_{\boldsymbol{A'}} \\ &- \sum_{\substack{\boldsymbol{A} \in \mathbb{K}_1 \\ \boldsymbol{B} \in \mathbb{K}_{-1}}} q^{n_1 \boldsymbol{A}_1 + \boldsymbol{B}_2 m_1 + \boldsymbol{B}_2 \boldsymbol{A}_1} Q_{\boldsymbol{A}+\boldsymbol{B}+(3m_1-1,3n_1-1)} P_{\boldsymbol{A}} P_{\boldsymbol{B}} \end{aligned}$$

(1.5)

$$e^{(\mu)}_{12}(m_1,n_1) = Q_{(3m_1+1,3n_1+1)}$$

(1.6)

$$e^{(\mu)}_{11}(m_1,n_1) = \sum_{\boldsymbol{A} \in \mathbb{K}_1} q^{\boldsymbol{A}_1 n_1} Q_{(3m_1,3n_1)+\boldsymbol{A}} P_{\boldsymbol{A}} + \frac{1}{2} \mu \delta_{(m_1,n_1),(0,0)}$$

(1.7)

$$\begin{aligned} e^{(\mu)}_{22}(m_1,n_1) = &- \sum_{\boldsymbol{A} \in \mathbb{K}_1} q^{\boldsymbol{A}_2 m_1} Q_{(3m_1,3n_1)+\boldsymbol{A}} P_{\boldsymbol{A}} \\ &- \sum_{\boldsymbol{B} \in \mathbb{K}_{-1}} q^{\boldsymbol{B}_2 m_1} Q_{(3m_1,3n_1)+\boldsymbol{B}} P_{\boldsymbol{B}} - \frac{1}{2} \mu \delta_{(m_1,n_1),(0,0)} \end{aligned}$$

(1.8)

$$\begin{aligned} e_{23}^{(\mu)}(m_1,n_1) &= -q^{-m_1n_1}\mu P_{-(3m_1+1,3n_1+1)} \\ &\quad - \sum_{\substack{\boldsymbol{A}\in\mathbb{K}_1 \\ \boldsymbol{B}\in\mathbb{K}_{-1}}} q^{n_1\boldsymbol{B}_1+\boldsymbol{A}_2m_1+\boldsymbol{A}_2\boldsymbol{B}_1} Q_{\boldsymbol{A}+\boldsymbol{B}+(3m_1+1,3n_1+1)} P_{\boldsymbol{A}} P_{\boldsymbol{B}} \\ &\quad - \sum_{\boldsymbol{B},\boldsymbol{B}'\in\mathbb{K}_{-1}} q^{n_1\boldsymbol{B}'_1+\boldsymbol{B}_2m_1+\boldsymbol{B}_2\boldsymbol{B}'_1} Q_{\boldsymbol{B}+\boldsymbol{B}'+(3m_1+1,3n_1+1)} P_{\boldsymbol{B}} P_{\boldsymbol{B}'} \end{aligned}$$

(1.9)

$$e_{32}^{(\mu)}(m_1,n_1) = Q_{(3m_1-1,3n_1-1)}$$

(1.10)

$$e_{31}^{(\mu)}(m_1,n_1) = \sum_{\boldsymbol{A}\in\mathbb{K}_1} q^{\boldsymbol{A}_1n_1} Q_{(3m_1-2,3n_1-2)+\boldsymbol{A}} P_{\boldsymbol{A}}$$

(1.11)

$$e_{13}^{(\mu)}(m_1,n_1) = \sum_{\boldsymbol{B}\in\mathbb{K}_{-1}} q^{\boldsymbol{B}_1n_1} Q_{(3m_1+2,3n_1+2)+\boldsymbol{B}} P_{\boldsymbol{B}}$$

(1.12)

$$e_{33}^{(\mu)}(m_1,n_1) = \sum_{\boldsymbol{B}\in\mathbb{K}_{-1}} q^{\boldsymbol{B}_1n_1} Q_{(3m_1,3n_1)+\boldsymbol{B}} P_{\boldsymbol{B}} + \frac{1}{2}\mu\delta_{(m_1,n_1),(0,0)}$$

$$D_1^{(\mu)} = \sum_{\boldsymbol{A}\in\mathbb{K}_1} \boldsymbol{A}_1 Q_{\boldsymbol{A}} P_{\boldsymbol{A}} + \sum_{\boldsymbol{B}\in\mathbb{K}_{-1}} \boldsymbol{B}_1 Q_{\boldsymbol{B}} P_{\boldsymbol{B}} \tag{1.13}$$

$$D_2^{(\mu)} = \sum_{\boldsymbol{A}\in\mathbb{K}_1} \boldsymbol{A}_2 Q_{\boldsymbol{A}} P_{\boldsymbol{A}} + \sum_{\boldsymbol{B}\in\mathbb{K}_{-1}} \boldsymbol{B}_2 Q_{\boldsymbol{B}} P_{\boldsymbol{B}} \tag{1.14}$$

Although the operators are infinite sums, they are well-defined as operators on V. Now we have the following result:

Theorem 1.15. *The linear map* $\pi : \widetilde{\mathfrak{gl}_3(\mathbb{C}_q)} \to$ *End* V *given by*

$$\begin{aligned} &\pi(E_{ij}(s^{m_1}t^{n_1})) = e_{ij}^{(\mu)}(m_1,n_1), \\ &\pi(d_s) = D_1^{(\mu)}, \quad \pi(d_t) = D_2^{(\mu)}, \quad \pi(c_s) = \pi(c_t) = 0, \end{aligned}$$

for $m_1, n_1 \in \mathbb{Z}, 1 \le i, j \le 3$, *is a Lie algebra homomorphism.*

Proof. The proof is straightforward. However, we would like to provide a few details. We shall do this orderly so that we won't miss any case.

First, we have

$$\begin{aligned}
&[e_{11}^{(\mu)}(m_1,n_1), e_{11}^{(\mu)}(m_2,n_2)]\\
=&[\sum_{\boldsymbol{A}\in\mathbb{K}_1} q^{\boldsymbol{A}_1 n_1} Q_{(3m_1,3n_1)+\boldsymbol{A}} P_{\boldsymbol{A}}, \sum_{\boldsymbol{A}'\in\mathbb{K}_1} q^{\boldsymbol{A}'_1 n_1} Q_{(3m_2,3n_2)+\boldsymbol{A}'} P_{\boldsymbol{A}'}]\\
=&\sum_{\boldsymbol{A}'\in\mathbb{K}_1} q^{(m_2+\boldsymbol{A}'_1)n_1+\boldsymbol{A}'n_2} Q_{(3m_1,3n_1)+(3m_2+3n_2)+\boldsymbol{A}'} P_{\boldsymbol{A}'} + \frac{1}{2}q^{m_2n_1}\mu\delta_{(m_1+m_2,n_1+n_2),(0,0)}\\
&-\sum_{\boldsymbol{A}\in\mathbb{K}_1} q^{\boldsymbol{A}_1 n_1+(m_1+\boldsymbol{A}_1)n_2} q_{(3m_1,3n_1)+(3m_2,3n_2)+\boldsymbol{A}} P_{\boldsymbol{A}} - \frac{1}{2}q^{m_1n_2}\mu\delta_{(m_1+m_2,n_1+n_2),(0,0)}\\
=&q^{m_2n_1}e_{11}^{(\mu)}(m_1+m_2,n_1+n_2) - q^{m_1n_2}e_{11}^{(\mu)}(m_1+m_2,n_1+n_2).
\end{aligned}$$

The following two brackets are easy.

$$\begin{aligned}
[e_{11}^{(\mu)}(m_1,n_1), e_{12}^{(\mu)}(m_2,n_2)] &= q^{m_2n_1}e_{12}^{(\mu)}(m_1+m_2,n_1+n_2)\\
[e_{11}^{(\mu)}(m_1,n_1), e_{13}^{(\mu)}(m_2,n_2)] &= q^{m_2n_1}e_{13}^{(\mu)}(m_1+m_2,n_1+n_2).
\end{aligned}$$

$$\begin{aligned}
&[e_{11}^{(\mu)}(m_1,n_1), e_{21}^{(\mu)}(m_2,n_2)]\\
=&\sum_{\boldsymbol{A}\in\mathbb{K}_1}(-\mu)q^{\boldsymbol{A}_1n_1-m_2n_2}[Q_{(3m_1,3n_1)+\boldsymbol{A}}P_{\boldsymbol{A}}, P_{-(3m_2-1,3n_2-1)}]\\
&-\sum_{\substack{\boldsymbol{A}\in\mathbb{K}_1\\ \bar{\boldsymbol{A}},\bar{\boldsymbol{A}}'\in\mathbb{K}_1}} q^{\boldsymbol{A}_1n_1+\bar{\boldsymbol{A}}'_1n_2+\bar{\boldsymbol{A}}_2m_2+\bar{\boldsymbol{A}}_2\bar{\boldsymbol{A}}'_1}[Q_{(3m_1,3n_1)+\boldsymbol{A}}P_{\boldsymbol{A}}, Q_{\bar{\boldsymbol{A}}+\bar{\boldsymbol{A}}'+(3m_2-1,3n_2-1)}P_{\bar{\boldsymbol{A}}}P_{\bar{\boldsymbol{A}}'}]\\
&-\sum_{\substack{\boldsymbol{A}\in\mathbb{K}_1\\ \bar{\boldsymbol{A}}\in\mathbb{K}_1\\ \boldsymbol{B}\in\mathbb{K}_{-1}}} q^{\boldsymbol{A}_1n_1+n_2\bar{\boldsymbol{A}}_1+\boldsymbol{B}_2m_2+\boldsymbol{B}_2\bar{\boldsymbol{A}}_1}[Q_{(3m_1,3n_1)+\boldsymbol{A}}P_{\boldsymbol{A}}, Q_{\bar{\boldsymbol{A}}+\boldsymbol{B}+(3m_2-1,3n_2-1)}P_{\bar{\boldsymbol{A}}}P_{\boldsymbol{B}}]\\
=&\mu q^{-(m_1+m_2)(n_1+n_2)+m_1n_2}P_{-(3(m_1+m_2)-1,3(n_1+n_2)-1)}\\
&-\sum_{\bar{\boldsymbol{A}},\bar{\boldsymbol{A}}'\in\mathbb{K}_1} q^{(\bar{\boldsymbol{A}}_1+\bar{\boldsymbol{A}}'_1+m_2)n_1+n_2\bar{\boldsymbol{A}}'_1+\bar{\boldsymbol{A}}_2m_2+\bar{\boldsymbol{A}}_2\bar{\boldsymbol{A}}'_1}Q_{(3m_1,3n_1)+\bar{\boldsymbol{A}}+\bar{\boldsymbol{A}}'+(3m_2-1,3n_2-1)}P_{\bar{\boldsymbol{A}}}P_{\bar{\boldsymbol{A}}'}\\
&+\sum_{\boldsymbol{A},\bar{\boldsymbol{A}}\in\mathbb{K}_1} q^{\boldsymbol{A}_1n_1+n_2(m_1+\boldsymbol{A}_1)+\bar{\boldsymbol{A}}_2m_2+\bar{\boldsymbol{A}}_2(m_1+\boldsymbol{A}_1)}Q_{\bar{\boldsymbol{A}}+(3m_1,3n_1)+\boldsymbol{A}+(3m_2-1,3n_2-1)}P_{\bar{\boldsymbol{A}}}P_{\boldsymbol{A}}\\
&+\sum_{\boldsymbol{A},\bar{\boldsymbol{A}}'\in\mathbb{K}_1} q^{\boldsymbol{A}_1n_1+n_2\boldsymbol{A}'_1+(n_1+\boldsymbol{A}_2)m_2+(n_1+\boldsymbol{A}_2)\bar{\boldsymbol{A}}'_1}Q_{\boldsymbol{A}+(3m_1,3n_1)+\bar{\boldsymbol{A}}'+(3m_2-1,3n_2-1)}P_{\bar{\boldsymbol{A}}'}P_{\boldsymbol{A}}\\
&+\sum_{\substack{\boldsymbol{A}\in\mathbb{K}_1\\ \boldsymbol{B}\in\mathbb{K}_{-1}}} q^{\boldsymbol{A}_1n_1+n_2(m_1+\boldsymbol{A}_1)+bdb_2m_2+\boldsymbol{B}_2(m_1+\boldsymbol{A}_1)}Q_{\boldsymbol{A}+(3m_1,3n_1)+\boldsymbol{B}+(3m_2-1,3n_2-1)}P_{\boldsymbol{B}}P_{\boldsymbol{A}}
\end{aligned}$$

(the second term and the fourth term are negative to each other)

$$
\begin{aligned}
=&-q^{m_1n_2}(-\mu q^{-(m_1+m_2)(n_1+n_2)}P_{-(3(m_1+m_2)-1,3(n_1+n_2)-1)}\\
&-\sum_{\boldsymbol{A},\bar{\boldsymbol{A}}\in\mathbb{K}_1} q^{\boldsymbol{A}(n_1+n_2)+\bar{\boldsymbol{A}}_2(m_1+m_2)+\boldsymbol{A}_1\bar{\boldsymbol{A}}_2}Q_{(3(m_1+m_2)-1,3(n_1+n_2)-1)+\boldsymbol{A}+\bar{\boldsymbol{A}}}P_{\bar{\boldsymbol{A}}}P_{\boldsymbol{A}}\\
&-\sum_{\substack{\boldsymbol{A}\in\mathbb{K}_1\\ \boldsymbol{B}\in\mathbb{K}_{-1}}} q^{\boldsymbol{A}_1(n_1+n_2)+\boldsymbol{B}_2(m_1+m_2)+\boldsymbol{B}_2+\boldsymbol{A}_1}Q_{\boldsymbol{A}+\boldsymbol{B}+(3(m_1+m_2)-1,3(n_1+n_2)-1)}P_{\boldsymbol{B}}P_{\boldsymbol{A}})\\
=&-q^{m_1n_2}e_{21}^{(\mu)}(m_1+m_2,n_1+n_2).
\end{aligned}
$$

The following seven brackets are easily verified.

$$
\begin{aligned}
&[e_{11}^{(\mu)}(m_1,n_1),e_{22}^{(\mu)}(m_2,n_2)]=0\\
&[e_{11}^{(\mu)}(m_1,n_1),e_{23}^{(\mu)}(m_2,n_2)]=0\\
&[e_{11}^{(\mu)}(m_1,n_1),e_{31}^{(\mu)}(m_2,n_2)]=-q^{m_1n_2}e_{31}^{(\mu)}(m_1+m_2,n_1+n_2)\\
&[e_{11}^{(\mu)}(m_1,n_1),e_{32}^{(\mu)}(m_2,n_2)]=0\\
&[e_{11}^{(\mu)}(m_1,n_1),e_{33}^{(\mu)}(m_2,n_2)]=0\\
&[e_{12}^{(\mu)}(m_1,n_1),e_{12}^{(\mu)}(m_2,n_2)]=0\\
&[e_{12}^{(\mu)}(m_1,n_1),e_{13}^{(\mu)}(m_2,n_2)]=0.
\end{aligned}
$$

Next, we have

$$
\begin{aligned}
&[e_{12}^{(\mu)}(m_1,n_1),e_{21}^{(\mu)}(m_2,n_2)]\\
=&\mu q^{-m_2n_2}\delta_{(m_1,n_1),(-m_2,n_2)}\\
&+\sum_{\boldsymbol{A}'\in\mathbb{K}_1} q^{n_2\boldsymbol{A}_1'+n_1m_2+n_1\boldsymbol{A}_1'}Q_{(3m_1+1,3n_1+1)+\boldsymbol{A}'+(3m_2-1,3n_2-1)}P_{\boldsymbol{A}'}\\
&+\sum_{\boldsymbol{A}\in\mathbb{K}_1} q^{n_2m_1+\boldsymbol{A}_2m_2+\boldsymbol{A}_2m_1}Q_{\boldsymbol{A}+(3m_1+1,3n_1+1)+(3m_2-1,3n_2-1)}P_{\boldsymbol{A}}\\
&+\sum_{\boldsymbol{B}\in\mathbb{K}_{-1}} q^{n_2m_1+\boldsymbol{B}_2m_2+\boldsymbol{B}_2m_1}Q_{(3m_1+1,3n_1+1)+\boldsymbol{B}+(3m_2-1,3n_2-1)}P_{\boldsymbol{B}}
\end{aligned}
$$

$$
\begin{aligned}
=&q^{n_1m_2}(\sum_{\boldsymbol{A'}\in\mathbb{K}_1} q^{(n_1+n_2)\boldsymbol{A}'_1}Q_{(3m_1+3m_2,3n_1+3n_2)+\boldsymbol{A'}}P_{\boldsymbol{A'}}+\frac{1}{2}\delta_{(m_1,n_1),(-m_2,n_2)})\\
&-q^{n_2m_1}(-\sum_{\boldsymbol{A}\in\mathbb{K}_1} q^{\boldsymbol{A}_2(m_2+m_1)}Q_{\boldsymbol{A}+(3m_1+3m_2,3n_1+3n_2)}P_{\boldsymbol{A}}\\
&-\sum_{\boldsymbol{B}\in\mathbb{K}_{-1}} q^{\boldsymbol{B}_2(m_2+m_1)}Q_{(3m_1+3m_2,3n_1+3n_2)+\boldsymbol{B}}P_{\boldsymbol{B}}-\frac{1}{2}\delta_{(m_1,n_1),(-m_2,n_2)})\\
=&q^{n_1m_2}e_{11}^{(\mu)}(m_1+m_2,n_1+n_2)-q^{n_2m_1}e_{22}^{(\mu)}(m_1+m_2,n_1+n_2).
\end{aligned}
$$

The following six brackets can be checked easily.

$$
\begin{aligned}
&[e_{12}^{(\mu)}(m_1,n_1),e_{22}^{(\mu)}(m_2,n_2)]=q^{n_1m_2}e_{12}^{(\mu)}(m_1+m_2,n_1+n_2)\\
&[e_{12}^{(\mu)}(m_1,n_1),e_{23}^{(\mu)}(m_2,n_2)]=q^{n_1m_2}e_{13}^{(\mu)}(m_1+m_2,n_1+n_2)\\
&[e_{12}^{(\mu)}(m_1,n_1),e_{31}^{(\mu)}(m_2,n_2)]=-q^{m_1n_2}e_{32}^{(\mu)}(m_1+m_2,n_1+n_2)\\
&[e_{12}^{(\mu)}(m_1,n_1),e_{32}^{(\mu)}(m_2,n_2)]=0\\
&[e_{12}^{(\mu)}(m_1,n_1),e_{33}^{(\mu)}(m_2,n_2)]=0\\
&[e_{13}^{(\mu)}(m_1,n_1),e_{13}^{(\mu)}(m_2,n_2)]=0.
\end{aligned}
$$

$$
\begin{aligned}
&[e_{13}^{(\mu)}(m_1,n_1),e_{21}^{(\mu)}(m_2,n_2)]\\
=&[\sum_{\boldsymbol{B}\in\mathbb{K}_{-1}} q^{\boldsymbol{B}_1n_1}Q_{(3m_1+2,3n_1+2)+\boldsymbol{B}}P_{\boldsymbol{B}},-q^{-m_2n_2}\mu P_{-(3m_2-1,3n_2-1)}\\
&-\sum_{\boldsymbol{A},bda'\in\mathbb{K}_1} q^{n_2\boldsymbol{A}'_1+\boldsymbol{A}_2m_2+\boldsymbol{A}_2\boldsymbol{A}'_1}Q_{\boldsymbol{A}+\boldsymbol{A'}+(3m_2-1,3n_2-1)}P_{\boldsymbol{A}}P_{\boldsymbol{A'}}\\
&-\sum_{\substack{\boldsymbol{A}\in\mathbb{K}_1\\ \boldsymbol{B}\in\mathbb{K}_{-1}}} q^{n_2\boldsymbol{A}_1+\boldsymbol{B}_2m_2+\boldsymbol{B}_2\boldsymbol{A}_1}Q_{\boldsymbol{A}+\boldsymbol{B}+(3m_2-1,3n_2-1)}P_{\boldsymbol{A}}P_{\boldsymbol{B}}]
\end{aligned}
$$

$$\begin{aligned}
=&-\sum_{\substack{\boldsymbol{A}\in\mathbb{K}_1\\ \boldsymbol{B}\in\mathbb{K}_{-1}}} q^{(\boldsymbol{A}_1+\boldsymbol{B}_1+m_2)n_1+n_2+\boldsymbol{A}_1+\boldsymbol{B}_2m_2+\boldsymbol{B}_2\boldsymbol{A}_1} Q_{(3m_1+2,3n_1+2)+\boldsymbol{A}+\boldsymbol{B}+(3m_2-1,3n_2-1)}P_{\boldsymbol{A}}P_{\boldsymbol{B}}\\
&+q^{-m_2n_2+(-m_1-m_2)n_1}\mu P_{(-3m_1-3m_2-1,-3n_1-3n_2-1)}\\
&+\sum_{\substack{\boldsymbol{A}\in\mathbb{K}_1\\ \boldsymbol{B}\in\mathbb{K}_{-1}}} q^{n_2(m_1+\boldsymbol{B}_1)+\boldsymbol{A}_2m_2+\boldsymbol{A}_2(m_1+\boldsymbol{B}_1)+\boldsymbol{B}_1n_1} Q_{\boldsymbol{A}+(3m_1+2,3n_1+2)+\boldsymbol{B}+(3m_2-1,3n_2-1)}P_{\boldsymbol{A}}P_{\boldsymbol{B}}\\
&+\sum_{\substack{\boldsymbol{A}'\in\mathbb{K}_1\\ \boldsymbol{B}\in\mathbb{K}_{-1}}} q^{n_2\boldsymbol{A}'_1+(n_1+\boldsymbol{B}_2)m_2+(n_1+\boldsymbol{B}_2)\boldsymbol{A}'_1+\boldsymbol{B}_1n_1} Q_{\boldsymbol{A}'+(3m_1+2,3n_1+2)+\boldsymbol{B}+(3m_2-1,3n_2-1)}P_{\boldsymbol{A}'}P_{\boldsymbol{B}}\\
&+\sum_{\boldsymbol{B},\boldsymbol{B}'\in\mathbb{K}_{-1}} q^{n_2(\boldsymbol{B}'_1+m_1)+\boldsymbol{B}_2m_2+\boldsymbol{B}_2(\boldsymbol{B}'_1+m_1)+\boldsymbol{B}'_1n_1} Q_{\boldsymbol{B}'+(3m_1+2,3n_1+2)+\boldsymbol{B}+(3m_2-1,3n_2-1)}P_{\boldsymbol{B}'}P_{\boldsymbol{B}}
\end{aligned}$$

(the first and the fourth terms cancel each other)

$$\begin{aligned}
=&q^{n_2m_1}(q^{-(m_1+m_2)(n_1+n_2)}\mu P_{-(3m_1+3m_2-1,3n_1+3n_2-1)}\\
&+\sum_{\boldsymbol{A},bda'\in\mathbb{K}_1} q^{(n_1+n_2)\boldsymbol{A}'_1+\boldsymbol{A}_2(m_1+m_2)+\boldsymbol{A}_2\boldsymbol{A}'_1} Q_{\boldsymbol{A}+\boldsymbol{A}'+(3m_1+3m_2-1,3n_1+3n_2-1)}P_{\boldsymbol{A}}P_{\boldsymbol{A}'}\\
&-\sum_{\substack{\boldsymbol{A}\in\mathbb{K}_1\\ \boldsymbol{B}\in\mathbb{K}_{-1}}} q^{(n_1+n_2)\boldsymbol{A}_1+\boldsymbol{B}_2(m_1+m_2)+\boldsymbol{B}_2\boldsymbol{A}_1} Q_{\boldsymbol{A}+\boldsymbol{B}+(3m_1+3m_2-1,3n_1+3n_2-1)}P_{\boldsymbol{A}}P_{\boldsymbol{B}})\\
=&-q^{n_2m_1}e^{(\mu)}_{23}(m_1+m_2,n_1+n_2).
\end{aligned}$$

The following two brackets are easy.

$$\begin{aligned}
[e^{(\mu)}_{13}(m_1,n_1),e^{(\mu)}_{22}(m_2,n_2)]&=0\\
[e^{(\mu)}_{13}(m_1,n_1),e^{(\mu)}_{23}(m_2,n_2)]&=0.
\end{aligned}$$

$$\begin{aligned}
&-[e^{(\mu)}_{13}(m_2,n_2),e^{(\mu)}_{31}(m_1,n_1)]=[e^{(\mu)}_{31}(m_1,n_1),e^{(\mu)}_{13}(m_2,n_2)]\\
=&\sum_{\substack{\boldsymbol{A}\in\mathbb{K}_1\\ \boldsymbol{B}\in\mathbb{K}_{-1}}} q^{n_1\boldsymbol{A}_1+n_2\boldsymbol{B}_1}[Q_{(3m_1-2,3n_1-2)+\boldsymbol{A}}P_{\boldsymbol{A}},Q_{(3m_2+2,3n_2+2)+\boldsymbol{B}}P_{\boldsymbol{B}}]\\
=&\sum_{\boldsymbol{B}\in\mathbb{K}_{-1}} q^{n_2\boldsymbol{B}_1+n_1(m_2+\boldsymbol{B}_1)}Q_{(3m_1+3m_2,3n_1+3n_2)+\boldsymbol{B}}P_{\boldsymbol{B}}\\
&-\sum_{\boldsymbol{A}\in\mathbb{K}_1} q^{n_1\boldsymbol{A}_1+n_2(m_1+\boldsymbol{A}_1)}Q_{(3m_1+3m_2,3n_1+3n_2)+\boldsymbol{A}}P_{\boldsymbol{A}}
\end{aligned}$$

$$
\begin{aligned}
=&q^{n_1m_2}\Big(\sum_{\boldsymbol{B}\in\mathbb{K}_{-1}} q^{n_2\boldsymbol{B}_1+n_1(m_2+\boldsymbol{B}_1)}Q_{(3m_1+3m_2,3n_1+3n_2)+\boldsymbol{B}}P_{\boldsymbol{B}}+\frac{1}{2}\mu\delta_{(m_1+m_2,n_1+n_2),(0,0)}\Big)\\
&-q^{n_2m_1}\Big(\sum_{\boldsymbol{A}\in\mathbb{K}_{1}} q^{n_1\boldsymbol{A}_1+n_2(m_1+\boldsymbol{A}_1)}Q_{(3m_1+3m_2,3n_1+3n_2)+\boldsymbol{A}}P_{\boldsymbol{A}}+\frac{1}{2}\mu\delta_{(m_1+m_2,n_1+n_2),(0,0)}\Big)\\
=&q^{n_1m_2}e^{(\mu)}_{33}(m_1+m_2,n_1+n_2)-q^{n_2m_1}e^{(\mu)}_{11}(m_1+m_2,n_1+n_2).
\end{aligned}
$$

The following two brackets are easy.

$$
\begin{aligned}
[e^{(\mu)}_{13}(m_1,n_1),e^{(\mu)}_{32}(m_2,n_2)]&=q^{n_1m_2}e^{(\mu)}_{12}(m_1+m_2,n_1+n_2)\\
[e^{(\mu)}_{13}(m_1,n_1),e^{(\mu)}_{33}(m_2,n_2)]&=q^{n_1m_2}e^{(\mu)}_{13}(m_1+m_2,n_1+n_2).
\end{aligned}
$$

$$
\begin{aligned}
&[e^{(\mu)}_{21}(m_1,n_1),e^{(\mu)}_{21}(m_2,n_2)]\\
=&\mu q^{-m_1n_1}\sum_{\boldsymbol{A}\in\mathbb{K}_1} q^{(-m_1-m_2-\boldsymbol{A}_1)n_2+\boldsymbol{A}_2m_2+\boldsymbol{A}_2(-m_1-m_2-\boldsymbol{A}_1)}P_{\boldsymbol{A}}P_{(-3m_1-3m_2,-3n_1-3n_2)-\boldsymbol{A}}\\
&-\mu q^{-m_2n_2}\sum_{\boldsymbol{A}'\in\mathbb{K}_1} q^{n_1\boldsymbol{A}'_1+(-n_1-n_2-\boldsymbol{A}'_2)m_1+(-n_1-n_2-\boldsymbol{A}'_2)\boldsymbol{A}'_1}P_{\boldsymbol{A}'}P_{(-3m_1-3m_2,-3n_1-3n_2)-\boldsymbol{A}'}\\
&+\Big[\sum_{\boldsymbol{A},\boldsymbol{A}'\in\mathbb{K}_1} q^{n_1\boldsymbol{A}'_1+\boldsymbol{A}_2m_1+\boldsymbol{A}_2\boldsymbol{A}'_1}Q_{\boldsymbol{A}+\boldsymbol{A}'+(3m_1-1,3n_1-1)}P_{\boldsymbol{A}}P_{\boldsymbol{A}'},\\
&\sum_{\boldsymbol{A},\boldsymbol{A}'\in\mathbb{K}_1} q^{n_2\boldsymbol{A}'_1+\boldsymbol{A}_2m_2+\boldsymbol{A}_2\boldsymbol{A}'_1}Q_{\boldsymbol{A}+\boldsymbol{A}'+(3m_2-1,3n_2-1)}P_{\boldsymbol{A}}P_{\boldsymbol{A}'}\Big]\\
&+\Big[\sum_{\boldsymbol{A},\boldsymbol{A}'\in\mathbb{K}_1} q^{n_1\boldsymbol{A}'_1+\boldsymbol{A}_2m_1+\boldsymbol{A}_2\boldsymbol{A}'_1}Q_{\boldsymbol{A}+\boldsymbol{A}'+(3m_1-1,3n_1-1)}P_{\boldsymbol{A}}P_{\boldsymbol{A}'},\\
&\sum_{\substack{\boldsymbol{A}\in\mathbb{K}_1\\ \boldsymbol{B}\in\mathbb{K}_{-1}}} q^{n_2\boldsymbol{A}_1+\boldsymbol{B}_2m_2+\boldsymbol{B}_2\boldsymbol{A}_1}Q_{\boldsymbol{A}+\boldsymbol{B}+(3m_2-1,3n_2-1)}P_{\boldsymbol{A}}P_{\boldsymbol{B}}\Big]\\
&+\Big[\sum_{\substack{\boldsymbol{A}\in\mathbb{K}_1\\ \boldsymbol{B}\in\mathbb{K}_{-1}}} q^{n_1\boldsymbol{A}_1+\boldsymbol{B}_2m_1+\boldsymbol{B}_2\boldsymbol{A}_1}Q_{\boldsymbol{A}+\boldsymbol{B}+(3m_1-1,3n_1-1)}P_{\boldsymbol{A}}P_{\boldsymbol{B}},\\
&\sum_{\boldsymbol{A},\boldsymbol{A}'\in\mathbb{K}_1} q^{n_2\boldsymbol{A}'_1+\boldsymbol{A}_2m_2+\boldsymbol{A}_2\boldsymbol{A}'_1}Q_{\boldsymbol{A}+\boldsymbol{A}'+(3m_2-1,3n_2-1)}P_{\boldsymbol{A}}P_{\boldsymbol{A}'}\Big]\\
&+\Big[\sum_{\substack{\boldsymbol{A}\in\mathbb{K}_1\\ \boldsymbol{B}\in\mathbb{K}_{-1}}} q^{n_1\boldsymbol{A}_1+\boldsymbol{B}_2m_1+\boldsymbol{B}_2\boldsymbol{A}_1}Q_{\boldsymbol{A}+\boldsymbol{B}+(3m_1-1,3n_1-1)}P_{\boldsymbol{A}}P_{\boldsymbol{B}},\\
&\sum_{\substack{\boldsymbol{A}\in\mathbb{K}_1\\ \boldsymbol{B}\in\mathbb{K}_{-1}}} q^{n_2\boldsymbol{A}_1+\boldsymbol{B}_2m_2+\boldsymbol{B}_2\boldsymbol{A}_1}Q_{\boldsymbol{A}+\boldsymbol{B}+(3m_2-1,3n_2-1)}P_{\boldsymbol{A}}P_{\boldsymbol{B}}\Big]
\end{aligned}
$$

(the first and the second terms cancel each other)

$$
\begin{aligned}
&= \sum_{\boldsymbol{A},\bar{\boldsymbol{A}},\bar{\boldsymbol{A}}'\in\mathbb{K}_1} q^{n_1(\bar{\boldsymbol{A}}_1+\bar{\boldsymbol{A}}'_1+m_2)+\boldsymbol{A}_2m_1+\boldsymbol{A}_2(\bar{\boldsymbol{A}}_1+\bar{\boldsymbol{A}}'_1+m_2)+n_2\bar{\boldsymbol{A}}'_1+\bar{\boldsymbol{A}}_2m_2+\bar{\boldsymbol{A}}_2\bar{\boldsymbol{A}}'_1}\\
&\quad\cdot Q_{\boldsymbol{A}+\bar{\boldsymbol{A}}+\bar{\boldsymbol{A}}'+(3m_1-1,3n_1-1)+(3m_2-1,3n_2-1)}P_{\boldsymbol{A}}P_{\bar{\boldsymbol{A}}}P_{\bar{\boldsymbol{A}}'}\\
&\quad+\sum_{\boldsymbol{A}',\bar{\boldsymbol{A}},\bar{\boldsymbol{A}}'\in\mathbb{K}_1} q^{n_1\boldsymbol{A}'_1+(\bar{\boldsymbol{A}}_2+\bar{\boldsymbol{A}}'_2+n_2)m_1+(\bar{\boldsymbol{A}}_2+\bar{\boldsymbol{A}}'_2+n_2)\boldsymbol{A}'_1+n_2\bar{\boldsymbol{A}}'_1+\bar{\boldsymbol{A}}_2m_2+\bar{\boldsymbol{A}}_2\bar{\boldsymbol{A}}'_1}\\
&\quad\cdot Q_{\boldsymbol{A}'+\bar{\boldsymbol{A}}+\bar{\boldsymbol{A}}'+(3m_1-1,3n_1-1)+(3m_2-1,3n_2-1)}P_{\boldsymbol{A}'}P_{\bar{\boldsymbol{A}}}P_{\bar{\boldsymbol{A}}'}
\end{aligned}
$$

$$
\begin{aligned}
&-\sum_{\boldsymbol{A},\boldsymbol{A}',\bar{\boldsymbol{A}}\in\mathbb{K}_1} q^{n_2(\boldsymbol{A}_1+\boldsymbol{A}'_1+m_1)+\bar{\boldsymbol{A}}_2m_2+\bar{\boldsymbol{A}}_2(\boldsymbol{A}_1+\boldsymbol{A}'_1+m_1)+n_1\boldsymbol{A}'_1+\boldsymbol{A}_2m_1+\boldsymbol{A}_2\boldsymbol{A}'_1}\\
&\cdot Q_{\boldsymbol{A}+\boldsymbol{A}'+\bar{\boldsymbol{A}}+(3m_1-1,3n_1-1)+(3m_2-1,3n_2-1)}P_{\boldsymbol{A}}P_{\boldsymbol{A}'}P_{\bar{\boldsymbol{A}}}\\
&-\sum_{\boldsymbol{A},\boldsymbol{A}',\bar{\boldsymbol{A}}'\in\mathbb{K}_1} q^{n_2\bar{\boldsymbol{A}}'_1+(\boldsymbol{A}_2+\boldsymbol{A}'_2+n_1)m_2+(\boldsymbol{A}_2+\boldsymbol{A}'_2+n_1)\bar{\boldsymbol{A}}'_1+n_1\boldsymbol{A}'_1+\boldsymbol{A}_2m_1+\boldsymbol{A}_2\boldsymbol{A}'_1}\\
&\cdot Q_{\boldsymbol{A}+\boldsymbol{A}'+\bar{\boldsymbol{A}}'+(3m_1-1,3n_1-1)+(3m_2-1,3n_2-1)}P_{\boldsymbol{A}}P_{\boldsymbol{A}'}P_{\bar{\boldsymbol{A}}'}\\
&-\sum_{\substack{\boldsymbol{A},\boldsymbol{A}'\in\mathbb{K}_1\\ \boldsymbol{B}\in\mathbb{K}_{-1}}} q^{n_2(\boldsymbol{A}_1+\boldsymbol{A}'_1+m_1)+\boldsymbol{B}_2m_2+\boldsymbol{B}_2(\boldsymbol{A}_1+\boldsymbol{A}'_1+m_1)+n_1\boldsymbol{A}'_1+\boldsymbol{A}_2m_1\boldsymbol{A}_2\boldsymbol{A}'_1}\\
&\cdot Q_{\boldsymbol{A}+\boldsymbol{A}'+\boldsymbol{B}+(3m_1-1,3n_1-1)+(3m_2-1,3n_2-1)}P_{\boldsymbol{A}}P_{\boldsymbol{A}'}P_{\boldsymbol{B}}\\
&+\sum_{\substack{\boldsymbol{A},\boldsymbol{A}'\in\mathbb{K}_1\\ \boldsymbol{B}\in\mathbb{K}_{-1}}} q^{n_1(\boldsymbol{A}_1+\boldsymbol{A}'_1+m_2)+\boldsymbol{B}_2m_1+\boldsymbol{B}_2(\boldsymbol{A}_1+\boldsymbol{A}'_1+m_2)+n_2\boldsymbol{A}'_1+\boldsymbol{A}_2m_2+\boldsymbol{A}_2\boldsymbol{A}'_1}\\
&\cdot Q_{\boldsymbol{A}+\boldsymbol{A}'+\boldsymbol{B}+(3m_1-1,3n_1-1)+(3m_2-1,3n_2-1)}P_{\boldsymbol{A}}P_{\boldsymbol{A}'}P_{\boldsymbol{B}}\\
&+\sum_{\substack{\boldsymbol{A},\bar{\boldsymbol{A}}\in\mathbb{K}_1\\ \boldsymbol{B}\in\mathbb{K}_{-1}}} q^{n_1\boldsymbol{A}_1+(\bar{\boldsymbol{A}}_2+\boldsymbol{B}_2+n_2)m_1+(\bar{\boldsymbol{A}}_2+\boldsymbol{B}_2+n_2)\boldsymbol{A}_1+n_2\bar{\boldsymbol{A}}_1+\boldsymbol{B}_2m_2+\boldsymbol{B}_2\bar{\boldsymbol{A}}_1}\\
&\cdot Q_{\boldsymbol{A}+\bar{\boldsymbol{A}}+\boldsymbol{B}+(3m_1-1,3n_1-1)+(3m_2-1,3n_2-1)}P_{\boldsymbol{A}}P_{\bar{\boldsymbol{A}}}P_{\boldsymbol{B}}\\
&-\sum_{\substack{\boldsymbol{A},\bar{\boldsymbol{A}}\in\mathbb{K}_1\\ \bar{\boldsymbol{B}}\in\mathbb{K}_{-1}}} q^{n_2\boldsymbol{A}_1+(\bar{\boldsymbol{A}}_2+\bar{\boldsymbol{B}}_2+n_1)m_2+(\bar{\boldsymbol{A}}_2+\bar{\boldsymbol{B}}_2+n_1)\boldsymbol{A}_1+n_1\bar{\boldsymbol{A}}_1+\bar{\boldsymbol{B}}_2m_1+\bar{\boldsymbol{B}}_2\bar{\boldsymbol{A}}_1}\\
&\cdot Q_{\boldsymbol{A}+\bar{\boldsymbol{A}}+\bar{\boldsymbol{B}}+(3m_1-1,3n_1-1)+(3m_2-1,3n_2-1)}P_{\boldsymbol{A}}P_{\bar{\boldsymbol{A}}}P_{\bar{\boldsymbol{B}}}=0
\end{aligned}
$$

as the first term cancels the fourth, the second term cancels the third, the fifth term cancels the seventh, and the sixth term cancels the eighth.

$$
\begin{aligned}
&-[e_{21}^{(\mu)}(m_2,n_2),e_{22}^{(\mu)}(m_1,n_1)]=[e_{22}^{(\mu)}(m_1,n_1),e_{21}^{(\mu)}(m_2,n_2)]\\
=&\mu\sum_{\boldsymbol{A}\in\mathbb{K}_1} q^{\boldsymbol{A}_2 m_1-m_2n_2}[Q_{(3m_1,3n_1)+\boldsymbol{A}}P_{\boldsymbol{A}},P_{-(3m_2-1,3n_2-1)}]\\
&+\sum_{\boldsymbol{A},\boldsymbol{A}',\bar{\boldsymbol{A}}\in\mathbb{K}_1} q^{\boldsymbol{A}_2m_1+n_2\boldsymbol{A}'_1+\bar{\boldsymbol{A}}_2m_2+\bar{\boldsymbol{A}}_2\boldsymbol{A}'_1}[Q_{(3m_1,3n_1)+\boldsymbol{A}}P_{\boldsymbol{A}},Q_{\bar{\boldsymbol{A}}+\boldsymbol{A}'+(3m_2-1,3n_2-1)}P_{\boldsymbol{A}'}P_{\bar{\boldsymbol{A}}}]\\
&+\sum_{\substack{\boldsymbol{A},\bar{\boldsymbol{A}}\in\mathbb{K}_1\\ \boldsymbol{B}\in\mathbb{K}_{-1}}} q^{\boldsymbol{A}_2m_1+n_2\bar{\boldsymbol{A}}_1+\boldsymbol{B}_2m_2+\boldsymbol{B}_2\bar{\boldsymbol{A}}_1}[Q_{(3m_1,3n_1)+\boldsymbol{A}}P_{\boldsymbol{A}},Q_{\bar{\boldsymbol{A}}+\boldsymbol{B}+(3m_2-1,3n_2-1)}P_{\boldsymbol{A}}P_{\boldsymbol{B}}]\\
&+\sum_{\substack{\boldsymbol{A}\in\mathbb{K}_1\\ \boldsymbol{B},\bar{\boldsymbol{B}}\in\mathbb{K}_{-1}}} q^{\boldsymbol{B}_2m_1+n_2\boldsymbol{A}_1+\bar{\boldsymbol{B}}_2m_2+\bar{\boldsymbol{B}}_2\boldsymbol{A}_1}[Q_{(3m_1,3n_1)+\boldsymbol{B}}P_{\boldsymbol{B}},Q_{\boldsymbol{A}+\bar{\boldsymbol{B}}+(3m_2-1,3n_2-1)}P_{\boldsymbol{A}}P_{\bar{\boldsymbol{B}}}]\\
=&-\mu q^{-m_2n_2+(-n_1-n_2)m_1}P_{-3m_1-3m_2+1,-3n_1-3n_2+1}\\
&+\sum_{\boldsymbol{A},\boldsymbol{A}'\in\mathbb{K}_1} q^{(\boldsymbol{A}_2+\boldsymbol{A}'_2+n_2)m_1+n_2\boldsymbol{A}'_1+\boldsymbol{A}_2m_2+\boldsymbol{A}_2\boldsymbol{A}'_1}Q_{(3m_1,3n_1)+\boldsymbol{A}+\boldsymbol{A}'+(3m_2-1,3n_2-1)}P_{\boldsymbol{A}}P_{\boldsymbol{A}'}\\
&-\sum_{\boldsymbol{A},\bar{\boldsymbol{A}}\in\mathbb{K}_1} q^{n_2(m_1+\bar{\boldsymbol{A}}_1)+\boldsymbol{A}_2m_2+\boldsymbol{A}_2(m_1+\bar{\boldsymbol{A}}_1)+\bar{\boldsymbol{A}}_2m_1}Q_{\boldsymbol{A}+\bar{\boldsymbol{A}}+(3m_1,3n_1)+(3m_2-1,3n_2-1)}P_{\boldsymbol{A}}P_{\bar{\boldsymbol{A}}}\\
&-\sum_{\boldsymbol{A}',\bar{\boldsymbol{A}}\in\mathbb{K}_1} q^{n_2\boldsymbol{A}'_1+(\bar{\boldsymbol{A}}_2+n_1)m_2+(\bar{\boldsymbol{A}}_2+n_1)\boldsymbol{A}'_1+\bar{\boldsymbol{A}}_2m_1}Q_{\boldsymbol{A}'+\bar{\boldsymbol{A}}+(3m_1,3n_1)+(3m_2-1,3n_2-1)}P_{\boldsymbol{A}'}P_{\bar{\boldsymbol{A}}}\\
&-\sum_{\substack{\boldsymbol{A}\in\mathbb{K}_1\\ \boldsymbol{B}\in\mathbb{K}_{-1}}} q^{n_2(\boldsymbol{A}_1+m_1)+\boldsymbol{B}_2m_2+\boldsymbol{B}_2(\boldsymbol{A}_1+m_1)+\boldsymbol{A}_2m_1}Q_{\boldsymbol{A}+\boldsymbol{B}+(3m_1,3n_1)+(3m_2-1,3n_2-1)}P_{\boldsymbol{A}}P_{\boldsymbol{B}}\\
&+\sum_{\substack{\boldsymbol{A}\in\mathbb{K}_1\\ \boldsymbol{B}\in\mathbb{K}_{-1}}} q^{(\boldsymbol{A}_2+\boldsymbol{B}_2+n_2)m_1+n_2\boldsymbol{A}_1+\boldsymbol{B}_2m_2+\boldsymbol{B}_2\boldsymbol{A}_1}Q_{\boldsymbol{A}+\boldsymbol{B}+(3m_1,3n_1)+(3m_2-1,3n_2-1)}P_{\boldsymbol{A}}P_{\boldsymbol{B}}\\
&-\sum_{\substack{\boldsymbol{A}\in\mathbb{K}_1\\ \boldsymbol{B}\in\mathbb{K}_{-1}}} q^{n_2\boldsymbol{A}_1+(n_1+\boldsymbol{B}_2)m_2+(n_1+\boldsymbol{B}_2)\boldsymbol{A}_1+\boldsymbol{B}_2m_1}Q_{\boldsymbol{A}+\boldsymbol{B}+(3m_1,3n_1)+(3m_2-1,3n_2-1)}P_{\boldsymbol{A}}P_{\boldsymbol{B}}
\end{aligned}
$$

(the second and the third terms are cancelled, and the fifth and the sixth terms are cancelled)

$$
\begin{aligned}
=&q^{n_1m_2}(-\mu q^{-(n_1+n_2)(m_1+m_2)}P_{-(3(m_1+m_2)-1,3(n_1+n_2)-1)}\\
&-\sum_{\bar{\boldsymbol{A}},\boldsymbol{A'}\in\mathbb{K}_1} q^{(n_1+n_2)\boldsymbol{A}'_1+(m_1+m_2)\bar{\boldsymbol{A}}_2+\bar{\boldsymbol{A}}_2\boldsymbol{A}'_1}Q_{(3m_1+3m_2-1,3n_1+3n_2-1)+\boldsymbol{A'}+\bar{\boldsymbol{A}}}P_{\boldsymbol{A'}}P_{\bar{\boldsymbol{A}}}\\
&-\sum_{\substack{\boldsymbol{A}\in\mathbb{K}_1\\ \boldsymbol{B}\in\mathbb{K}_{-1}}} q^{(n_1+n_2)\boldsymbol{A}_1+(m_1+m_2)\boldsymbol{B}_2+\boldsymbol{B}_2\boldsymbol{A}_1})Q_{(3m_1+3m_2-1,3n_1+3n_2-1)+\boldsymbol{A}+\boldsymbol{B}}P_{\boldsymbol{A}}P_{\boldsymbol{B}}\\
=&q^{n_1m_2}e_{21}^{(\mu)}(m_1+m_2,n_1+n_2).
\end{aligned}
$$

$$
\begin{aligned}
&[e_{21}^{(\mu)}(m_1,n_1),e_{23}^{(\mu)}(m_2,n_2)]\\
=&\mu q^{-m_1n_1}\sum_{\boldsymbol{A}\in\mathbb{K}_1} q^{n_2(-m_1-m_2-\boldsymbol{A}_1)+\boldsymbol{A}_2m_2+\boldsymbol{A}_2(-m_1-m_2-\boldsymbol{A}_1)}P_{\boldsymbol{A}}P_{(-3m_1-3m_2,-3n_1-3n_2)-\boldsymbol{A}}\\
&+[\sum_{\boldsymbol{A},\boldsymbol{A'}\in\mathbb{K}_1} q^{n_1\boldsymbol{A}'_1+\boldsymbol{A}_2m_1+\boldsymbol{A}_2\boldsymbol{A}'_1}Q_{\boldsymbol{A}+\boldsymbol{A'}+(3m_1-1,3n_1-1)}P_{\boldsymbol{A}}P_{\boldsymbol{A'}},\\
&\sum_{\substack{\boldsymbol{A}\in\mathbb{K}_1\\ \boldsymbol{B}\in\mathbb{K}_{-1}}} q^{n_2\boldsymbol{B}_1+\boldsymbol{A}_2m_2+\boldsymbol{A}_2\boldsymbol{B}_1}Q_{(3m_2+1,3n_2+1)+\boldsymbol{A}+\boldsymbol{B}}P_{\boldsymbol{A}}P_{\boldsymbol{B}}]\\
&-\mu q^{-m_2n_2}\sum_{\boldsymbol{A}\in\mathbb{K}_1} q^{n_1\boldsymbol{A}_1+(-n_1-n_2-\boldsymbol{A}_2)m_1+(-n_1-n_2-\boldsymbol{A}_2)\boldsymbol{A}_1}P_{\boldsymbol{A}}P_{(-3m_1-3m_2,-3n_1-3n_2)-\boldsymbol{A}}\\
&+[\sum_{\substack{\boldsymbol{A}\in\mathbb{K}_1\\ \boldsymbol{B}\in\mathbb{K}_{-1}}} q^{n_1\boldsymbol{A}_1+\boldsymbol{B}_2m_1+\boldsymbol{B}_2\boldsymbol{A}_1}Q_{\boldsymbol{A}+\boldsymbol{B}+(3m_1-1,3n_1-1)}P_{\boldsymbol{A}}P_{\boldsymbol{B}},\\
&\sum_{\substack{\boldsymbol{A}\in\mathbb{K}_1\\ \boldsymbol{B}\in\mathbb{K}_{-1}}} q^{n_2\boldsymbol{B}_1+\boldsymbol{A}_2m_2+\boldsymbol{A}_2\boldsymbol{B}_1}Q_{(3m_2+1,3n_2+1)}P_{\boldsymbol{A}}P_{\boldsymbol{B}}]\\
&+[\sum_{\substack{\boldsymbol{A}\in\mathbb{K}_1\\ \boldsymbol{B}\in\mathbb{K}_{-1}}} q^{n_1\boldsymbol{A}_1+\boldsymbol{B}_2m_1+\boldsymbol{B}_2\boldsymbol{A}_1}Q_{\boldsymbol{A}+\boldsymbol{B}+(3m_1-1,3n_1-1)}P_{\boldsymbol{A}}P_{\boldsymbol{B}},\\
&\sum_{\boldsymbol{B},\boldsymbol{B'}\in\mathbb{K}_{-1}} q^{n_2\boldsymbol{B}'_1+\boldsymbol{B}_2m_2+\boldsymbol{B}_2\boldsymbol{B}'_1}Q_{(3m_2+1,3n_2+1)+\boldsymbol{B}+\boldsymbol{B'}}P_{\boldsymbol{B}}P_{\boldsymbol{B'}}]
\end{aligned}
$$

(the first term and the third term are negative to each other)

$$
\begin{aligned}
= \sum_{\substack{\boldsymbol{A},\bar{\boldsymbol{A}}\in\mathbb{K}_1\\ \boldsymbol{B}\in\mathbb{K}_{-1}}} & q^{n_1(m_2+\bar{\boldsymbol{A}}_1+\boldsymbol{B}_1)+\boldsymbol{A}_2 m_1+\boldsymbol{A}_2(m_1+\bar{\boldsymbol{A}}_1+\boldsymbol{B}_1)+n_2\boldsymbol{B}_1+\bar{\boldsymbol{A}}_2 m_2+\bar{\boldsymbol{A}}_2\boldsymbol{B}_1} \\
&\cdot Q_{\boldsymbol{A}+\bar{\boldsymbol{A}}+\boldsymbol{B}+(3m_1+3m_2,3n_1+3n_2)} P_{\boldsymbol{A}} P_{\bar{\boldsymbol{A}}} P_{\boldsymbol{B}} \\
+ \sum_{\substack{\boldsymbol{A}',\bar{\boldsymbol{A}}\in\mathbb{K}_1\\ \boldsymbol{B}\in\mathbb{K}_{-1}}} & q^{n_1\boldsymbol{A}'_1+(n_2+\bar{\boldsymbol{A}}_2+\boldsymbol{B}_2)m_1+(n_2+\bar{\boldsymbol{A}}_2+\boldsymbol{B}_2)\bar{\boldsymbol{A}}_1+n_2\boldsymbol{B}_1+\bar{\boldsymbol{A}}_2 m_2+\bar{\boldsymbol{A}}_2\boldsymbol{B}_1} \\
&\cdot Q_{\boldsymbol{A}'+\bar{\boldsymbol{A}}+\boldsymbol{B}+(3m_1+3m_2,3n_1+3n_2)} P_{\boldsymbol{A}'} P_{\bar{\boldsymbol{A}}} P_{\boldsymbol{B}} \\
- \sum_{\substack{\boldsymbol{A},\boldsymbol{A}'\in\mathbb{K}_1\\ \boldsymbol{B}\in\mathbb{K}_{-1}}} & q^{n_2\boldsymbol{B}_1+(\boldsymbol{A}_2+\boldsymbol{A}'_2+n_1)m_2+(\boldsymbol{A}_2+\boldsymbol{A}'_2+n_1)\boldsymbol{B}_1+n_1\boldsymbol{A}'_1+\boldsymbol{A}_2 m_1+\boldsymbol{A}_2\boldsymbol{A}'_1} \\
&\cdot Q_{\boldsymbol{A}'+\boldsymbol{A}+\boldsymbol{B}+(3m_1+3m_2,3n_1+3n_2)} P_{\boldsymbol{A}'} P_{\boldsymbol{A}} P_{\boldsymbol{B}}
\end{aligned}
$$

$$
\begin{aligned}
+ \sum_{\substack{\boldsymbol{A}\in\mathbb{K}_1\\ \boldsymbol{B},\boldsymbol{B}'\in\mathbb{K}_{-1}}} & q^{n_1(m_2+\boldsymbol{A}_1+\boldsymbol{B}'_1)+\boldsymbol{B}_2 m_1+\boldsymbol{B}_2(m_2+\boldsymbol{A}_1+\boldsymbol{B}'_1)+n_2\boldsymbol{B}'_1+\boldsymbol{A}_2 m_2+\boldsymbol{A}_2\boldsymbol{B}'_1} \\
&\cdot Q_{\boldsymbol{A}+\boldsymbol{B}+\boldsymbol{B}'+(3m_1+3m_2,3n_1+3n_2)} P_{\boldsymbol{A}} P_{\boldsymbol{B}} P_{\boldsymbol{B}'} \\
- \sum_{\substack{\boldsymbol{A},\bar{\boldsymbol{A}}\in\mathbb{K}_1\\ \boldsymbol{B}\in\mathbb{K}_{-1}}} & q^{n_2(\boldsymbol{A}_1+\boldsymbol{B}_1+m_1)+\bar{\boldsymbol{A}}_2 m_2+\bar{\boldsymbol{A}}_2(\boldsymbol{A}_1+\boldsymbol{B}+m_1)+n_1\boldsymbol{A}_1+\boldsymbol{B}_2 m_1+\boldsymbol{B}_2\boldsymbol{A}_1} \\
&\cdot Q_{\boldsymbol{A}+\bar{\boldsymbol{A}}+\boldsymbol{B}+(3m_1+3m_2,3n_1+3n_2)} P_{\boldsymbol{A}} P_{\bar{\boldsymbol{A}}} P_{\boldsymbol{B}} \\
+ \sum_{\substack{\boldsymbol{A}\in\mathbb{K}_1\\ \boldsymbol{B},\boldsymbol{B}'\in\mathbb{K}_{-1}}} & q^{n_1\boldsymbol{A}_1+(n_2+\boldsymbol{B}_2+\boldsymbol{B}'_2)m_1+(n_2+\boldsymbol{B}_2+\boldsymbol{B}'_2)\boldsymbol{A}_1+n_2\boldsymbol{B}'_1+\boldsymbol{B}_2 m_2+\boldsymbol{B}_2\boldsymbol{B}'_1} \\
&\cdot Q_{\boldsymbol{A}+\boldsymbol{B}+\boldsymbol{B}'+(3m_1+3m_2,3n_1+3n_2)} P_{\boldsymbol{A}} P_{\boldsymbol{B}} P_{\boldsymbol{B}'} \\
- \sum_{\substack{\boldsymbol{A}\in\mathbb{K}_1\\ \boldsymbol{B},\bar{\boldsymbol{B}}\in\mathbb{K}_{-1}}} & q^{n_2(\boldsymbol{A}_1+\bar{\boldsymbol{B}}_1+m_1)+\boldsymbol{B}_2 m_2+\boldsymbol{B}_2(\boldsymbol{A}_1+\bar{\boldsymbol{B}}_1+m_1)+n_1\boldsymbol{A}_1+\bar{\boldsymbol{B}}_2 m_1+\bar{\boldsymbol{B}}_2\boldsymbol{A}_1} \\
&\cdot Q_{\boldsymbol{A}+\boldsymbol{B}+\bar{\boldsymbol{B}}+(3m_1+3m_2,3n_1+3n_2)} P_{\boldsymbol{A}} P_{\boldsymbol{B}} P_{\bar{\boldsymbol{B}}} \\
- \sum_{\substack{\boldsymbol{A}\in\mathbb{K}_1\\ \boldsymbol{B}',\bar{\boldsymbol{B}}\in\mathbb{K}_{-1}}} & q^{n_2\boldsymbol{B}'_1+(\boldsymbol{A}_2+\bar{\boldsymbol{B}}_2+n_1)m_2+(\boldsymbol{A}_2+\bar{\boldsymbol{B}}_2+n_1)\boldsymbol{B}'_1+n_1\boldsymbol{A}_1+\bar{\boldsymbol{B}}_2 m_1+\bar{\boldsymbol{B}}_2\boldsymbol{A}_1} \\
&\cdot Q_{\boldsymbol{A}+\boldsymbol{B}'+\bar{\boldsymbol{B}}+(3m_1+3m_2,3n_1+3n_2)} P_{\boldsymbol{A}} P_{\boldsymbol{B}'} P_{\bar{\boldsymbol{B}}} = 0
\end{aligned}
$$

as the first term and the third, the second and the fifth, the fourth and the eighth, the sixth and the seventh are cancelled.

The following three brackets are easy.

$$\begin{aligned}
&[e_{21}^{(\mu)}(m_1,n_1),e_{31}^{(\mu)}(m_2,n_2)]=0\\
&[e_{21}^{(\mu)}(m_1,n_1),e_{32}^{(\mu)}(m_2,n_2)]=-q^{n_2m_1}e_{31}^{(\mu)}(m_1+m_2,n_1+n_2)\\
&[e_{21}^{(\mu)}(m_1,n_1),e_{33}^{(\mu)}(m_2,n_2)]=0.
\end{aligned}$$

$$\begin{aligned}
&[e_{22}^{(\mu)}(m_1,n_1),e_{22}^{(\mu)}(m_2,n_2)]\\
=&\sum_{\boldsymbol{A},\boldsymbol{A'}\in\mathbb{K}_1}q^{\boldsymbol{A}_2m_1+\boldsymbol{A}'_2m_2}[Q_{(3m_1,3n_1)+\boldsymbol{A}}P_{\boldsymbol{A}},Q_{(3m_2,3n_2)+\boldsymbol{A'}}P_{\boldsymbol{A'}}]\\
&+\sum_{\boldsymbol{B},\boldsymbol{B'}\in\mathbb{K}_{-1}}q^{\boldsymbol{B}_2m_1+\boldsymbol{B}'_2m_2}[Q_{(3m_1,3n_1)+\boldsymbol{B}}P_{\boldsymbol{B}},Q_{(3m_2,3n_2)+\boldsymbol{B'}}P_{\boldsymbol{B'}}]\\
=&\sum_{\boldsymbol{A'}\in\mathbb{K}_1}q^{(n_2+\boldsymbol{A}'_2)m_1+\boldsymbol{A}'_2m_2}Q_{(3m_1,3n_1)+(3m_2,3n_2)+\boldsymbol{A'}}P_{\boldsymbol{A'}}\\
&-\sum_{\boldsymbol{A}\in\mathbb{K}_1}q^{\boldsymbol{A}_2m_1+(n_1+\boldsymbol{A}_2)m_2}Q_{(3m_1,3n_1)+(3m_2,3n_2)+\boldsymbol{A}}P_{\boldsymbol{A}}\\
&+\sum_{\boldsymbol{B'}\in\mathbb{K}_{-1}}q^{(n_2+\boldsymbol{B}'_2)m_1+\boldsymbol{B}'_2m_2}Q_{(3m_1,3n_1)+(3m_2,3n_2)+\boldsymbol{B'}}P_{\boldsymbol{B'}}\\
&-\sum_{\boldsymbol{B}\in\mathbb{K}_{-1}}q^{\boldsymbol{B}_2m_1+(n_1+\boldsymbol{B}_2)m_2}Q_{(3m_1,3n_1)+(3m_2,3n_2)+\boldsymbol{B}}P_{\boldsymbol{B}}\\
=&q^{n_1m_2}(-\sum_{\boldsymbol{A}\in\mathbb{K}_1}q^{\boldsymbol{A}_2(m_1+m_2)}Q_{(3m_1,3n_1)+(3m_2,3n_2)+\boldsymbol{A}}P_{\boldsymbol{A}}\\
&-\sum_{\boldsymbol{B}\in\mathbb{K}_{-1}}q^{\boldsymbol{B}_2(m_1+m_2)}Q_{(3m_1,3n_1)+(3m_2,3n_2)+\boldsymbol{B}}P_{\boldsymbol{B}}-\frac{1}{2}\mu\delta_{(m_1+m_2,n_1+n_2),(0,0)})\\
&-q^{n_2m_1}(-\sum_{\boldsymbol{A}\in\mathbb{K}_1}q^{\boldsymbol{A}_2(m_1+m_2)}Q_{(3m_1,3n_1)+(3m_2,3n_2)+\boldsymbol{A}}P_{\boldsymbol{A}}\\
&-\sum_{\boldsymbol{B}\in\mathbb{K}_{-1}}q^{\boldsymbol{B}_2(m_1+m_2)}Q_{(3m_1,3n_1)+(3m_2,3n_2)+\boldsymbol{B}}P_{\boldsymbol{B}}-\frac{1}{2}\mu\delta_{(m_1+m_2,n_1+n_2),(0,0)})\\
=&q^{n_1m_2}e_{22}^{(\mu)}(m_1+m_2,n_1+n_2)-q^{n_2m_1}e_{22}^{(\mu)}(m_1+m_2,n_1+n_2).
\end{aligned}$$

$$
\begin{aligned}
&[e_{22}^{(\mu)}(m_1,n_1), e_{23}^{(\mu)}(m_2,n_2)]\\
=&[-\sum_{\boldsymbol{A}\in\mathbb{K}_1} q^{\boldsymbol{A}_2 m_1} Q_{(3m_1,3n_1)+\boldsymbol{A}} P_{\boldsymbol{A}} - \sum_{\boldsymbol{B}\in\mathbb{K}_{-1}} q^{\boldsymbol{B}_2 m_1} Q_{(3m_1,3n_1)+\boldsymbol{B}} P_{\boldsymbol{B}},\\
&- q^{-m_2n_2}\mu P_{-(3m_2+1,3n_2+1)} - \sum_{\substack{\boldsymbol{A}\in\mathbb{K}_1\\ \boldsymbol{B}\in\mathbb{K}_{-1}}} q^{n_2\boldsymbol{B}_1+\boldsymbol{A}_2m_2+\boldsymbol{A}_2\boldsymbol{B}_1} Q_{\boldsymbol{A}+\boldsymbol{B}+(3m_2+1,3n_2+1)} P_{\boldsymbol{A}} P_{\boldsymbol{B}}\\
&- \sum_{\boldsymbol{B},\boldsymbol{B}'\in\mathbb{K}_{-1}} q^{n_2\boldsymbol{B}'_1+\boldsymbol{B}_2m_2+\boldsymbol{B}_2\boldsymbol{B}'_1} Q_{\boldsymbol{B}'+\boldsymbol{B}+(3m_2+1,3n_2+1)} P_{\boldsymbol{B}} P_{\boldsymbol{B}'}]\\
=&\sum_{\substack{\boldsymbol{A}\in\mathbb{K}_1\\ \boldsymbol{B}\in\mathbb{K}_{-1}}} q^{(n_2+\boldsymbol{A}_2+\boldsymbol{B}_2)m_1+n_2\boldsymbol{B}_1+\boldsymbol{A}_2m_2+\boldsymbol{A}_2\boldsymbol{B}_1} Q_{(3m_1,3n_1)+\boldsymbol{A}+\boldsymbol{B}+(3m_2+1,3n_2+1)} P_{\boldsymbol{A}} P_{\boldsymbol{B}}\\
&- \sum_{\substack{\boldsymbol{A}\in\mathbb{K}_1\\ \boldsymbol{B}\in\mathbb{K}_{-1}}} q^{n_2\boldsymbol{B}_1+(n_1+\boldsymbol{A}_2)m_2+(\boldsymbol{A}_2+n_1)\boldsymbol{B}_1+\boldsymbol{A}_2m_1} Q_{(3m_1,3n_1)+\boldsymbol{A}+\boldsymbol{B}+(3m_2+1,3n_2+1)} P_{\boldsymbol{A}} P_{\boldsymbol{B}}\\
&- q^{-m_2n_2+(-n_2-n_1)m_1}\mu P_{-(3m_2+1,3n_2+1)-(3m_1,3n_1)}\\
&- \sum_{\substack{\boldsymbol{A}\in\mathbb{K}_1\\ \boldsymbol{B}\in\mathbb{K}_{-1}}} q^{n_2(m_1+\boldsymbol{B}_1)+\boldsymbol{A}_2m_2+\boldsymbol{A}_2(m_1+\boldsymbol{B}_1)+\boldsymbol{B}_2m_1} Q_{(3m_1,3n_1)+\boldsymbol{A}+\boldsymbol{B}+(3m_2+1,3n_2+1)} P_{\boldsymbol{A}} P_{\boldsymbol{B}}\\
&+ \sum_{\boldsymbol{B},\boldsymbol{B}'\in\mathbb{K}_{-1}} q^{(n_2+\boldsymbol{B}_2+\boldsymbol{B}'_2)m_1+n_2\boldsymbol{B}'_1+\boldsymbol{B}_2m_2+\boldsymbol{B}_2\boldsymbol{B}'_1} Q_{(3m_1,3n_1)+\boldsymbol{B}+\boldsymbol{B}'+(3m_2+1,3n_2+1)} P_{\boldsymbol{B}} P_{\boldsymbol{B}'}\\
&- \sum_{\boldsymbol{B},\boldsymbol{B}'\in\mathbb{K}_{-1}} q^{n_2(\boldsymbol{B}'_1+m_1)+\boldsymbol{B}_2m_2+\boldsymbol{B}_2(\boldsymbol{B}'_1+m_1)+\boldsymbol{B}'_2m_1} Q_{(3m_1,3n_1)+\boldsymbol{B}+\boldsymbol{B}'+(3m_2+1,3n_2+1)} P_{\boldsymbol{B}} P_{\boldsymbol{B}'}\\
&- \sum_{\boldsymbol{B},\boldsymbol{B}'\in\mathbb{K}_{-1}} q^{n_2\boldsymbol{B}'_1+(n_1+\boldsymbol{B}_2)m_2+(\boldsymbol{B}_2+n_1)\boldsymbol{B}'_1+\boldsymbol{B}_2m_1} Q_{(3m_1,3n_1)+\boldsymbol{B}+\boldsymbol{B}'+(3m_2+1,3n_2+1)} P_{\boldsymbol{B}} P_{\boldsymbol{B}'}
\end{aligned}
$$

(the first term cancels the fourth while the fifth cancels the sixth)

$$
\begin{aligned}
=&q^{m_2n_1}(-q^{-(m_1+m_2)(n_1+n_2)}\mu P_{-(3m_1+3m_2+1,3n_1+3n_2+1)}\\
&- \sum_{\substack{\boldsymbol{A}\in\mathbb{K}_1\\ \boldsymbol{B}\in\mathbb{K}_{-1}}} q^{\boldsymbol{A}_2(m_1+m_2)+(n_1+n_2)\boldsymbol{B}_1+\boldsymbol{A}_2\boldsymbol{B}_1} Q_{\boldsymbol{A}+\boldsymbol{B}+(3m_1+3m_2+1,3n_1+3n_2+1)} P_{\boldsymbol{A}} P_{\boldsymbol{B}}\\
&- \sum_{\boldsymbol{B},\boldsymbol{B}'\in\mathbb{K}_{-1}} q^{\boldsymbol{B}'_1(n_1+n_2)+\boldsymbol{B}_2(m_1+m_2)\boldsymbol{B}_2\boldsymbol{B}'_1})Q_{\boldsymbol{B}+\boldsymbol{B}'+(3m_1+3m_2+1,3n_1+3n_2+1)} P_{\boldsymbol{B}} P_{\boldsymbol{B}'}\\
=&q^{m_2n_1} e_{23}^{(\mu)}(m_1+m_2, n_1+n_2).
\end{aligned}
$$

The following three brackets are easy.

$$
\begin{aligned}
&[e_{22}^{(\mu)}(m_1,n_1), e_{31}^{(\mu)}(m_2,n_2)] = 0\\
&[e_{22}^{(\mu)}(m_1,n_1), e_{32}^{(\mu)}(m_2,n_2)] = -q^{n_2m_1}e_{32}^{(\mu)}(m_1+m_2,n_1+n_2)\\
&[e_{22}^{(\mu)}(m_1,n_1), e_{33}^{(\mu)}(m_2,n_2)] = 0.
\end{aligned}
$$

$$
\begin{aligned}
&[e_{23}^{(\mu)}(m_1,n_1), e_{23}^{(\mu)}(m_2,n_2)]\\
=&[-q^{-m_1n_1}\mu P_{-(3m_1+1,3n_1+1)} - \sum_{\substack{\boldsymbol{A}\in\mathbb{K}_1\\ \boldsymbol{B}\in\mathbb{K}_{-1}}} q^{n_1\boldsymbol{B}_1+\boldsymbol{A}_2m_1+\boldsymbol{A}_2\boldsymbol{B}_1}Q_{\boldsymbol{A}+\boldsymbol{B}+(3m_1+1,3n_1+1)}P_{\boldsymbol{A}}P_{\boldsymbol{B}}\\
&-\sum_{\boldsymbol{B},\boldsymbol{B}'\in\mathbb{K}_{-1}} q^{n_1\boldsymbol{B}'_1+\boldsymbol{B}_2m_1+\boldsymbol{B}_2\boldsymbol{B}'_1}Q_{\boldsymbol{B}+\boldsymbol{B}'+(3m_1+1,3n_1+1)}P_{\boldsymbol{B}}P_{\boldsymbol{B}'},\\
&-q^{-m_2n_2}\mu P_{-(3m_2+1,3n_2+1)} - \sum_{\substack{\boldsymbol{A}\in\mathbb{K}_1\\ \boldsymbol{B}\in\mathbb{K}_{-1}}} q^{n_2\boldsymbol{B}_1+\boldsymbol{A}_2m_2+\boldsymbol{A}_2\boldsymbol{B}_1}Q_{\boldsymbol{A}+\boldsymbol{B}+(3m_2+1,3n_2+1)}P_{\boldsymbol{A}}P_{\boldsymbol{B}}\\
&-\sum_{\boldsymbol{B},\boldsymbol{B}'\in\mathbb{K}_{-1}} q^{n_2\boldsymbol{B}'_1+\boldsymbol{B}_2m_2+\boldsymbol{B}_2\boldsymbol{B}'_1}Q_{\boldsymbol{B}+\boldsymbol{B}'+(3m_2+1,3n_2+1)}P_{\boldsymbol{B}}P_{\boldsymbol{B}'}]\\
=&\sum_{\boldsymbol{B}\in\mathbb{K}_{-1}} \mu q^{-m_1n_1+n_2(-m_1-m_2-\boldsymbol{B}_1)+\boldsymbol{B}_2m_2+\boldsymbol{B}_2(-m_1-m_2-\boldsymbol{B}_1)}P_{\boldsymbol{B}}P_{-(3m_1+1,3n_1+1)-(3m_2+1,3n_2+1)-\boldsymbol{B}}\\
&-\sum_{\boldsymbol{B}\in\mathbb{K}_{-1}} \mu q^{-m_2n_2+n_1\boldsymbol{B}_1+(-n_2-n_1-\boldsymbol{B}_2)m_1+(-n_2-n_1-\boldsymbol{B}_2)\boldsymbol{B}_1}P_{\boldsymbol{B}}P_{-(3m_1+1,3n_1+1)-(3m_2+1,3n_2+1)-\boldsymbol{B}}\\
&+\sum_{\boldsymbol{B},\bar{\boldsymbol{B}},\boldsymbol{B}'\in\mathbb{K}_{-1}} q^{n_1(\boldsymbol{B}'_1+\bar{\boldsymbol{B}}_1+m_2)+\boldsymbol{B}_2m_1+\boldsymbol{B}_2(\boldsymbol{B}'_1+\bar{\boldsymbol{B}}_1+m_2)+n_2\boldsymbol{B}'_1+\bar{\boldsymbol{B}}_2m_2+\bar{\boldsymbol{B}}_2\boldsymbol{B}'_1}\\
&\cdot Q_{\boldsymbol{B}+\boldsymbol{B}'+\bar{\boldsymbol{B}}+(3m_1+1,3n_1+1)+(3m_2+1,3n_2+1)}P_{\boldsymbol{B}}P_{\boldsymbol{B}'}P_{\bar{\boldsymbol{B}}}\\
&+\sum_{\boldsymbol{B},\bar{\boldsymbol{B}},\boldsymbol{B}'\in\mathbb{K}_{-1}} q^{n_1\boldsymbol{B}'_1+(\boldsymbol{B}_2+\bar{\boldsymbol{B}}_2+n_2)m_1+\boldsymbol{B}'_1(\boldsymbol{B}_2+\bar{\boldsymbol{B}}_2+n_2)+n_2\bar{\boldsymbol{B}}_1+\boldsymbol{B}_2m_2+\boldsymbol{B}_2\bar{\boldsymbol{B}}_1}\\
&\cdot Q_{\boldsymbol{B}+\boldsymbol{B}'+\bar{\boldsymbol{B}}+(3m_1+1,3n_1+1)+(3m_2+1,3n_2+1)}P_{\boldsymbol{B}}P_{\boldsymbol{B}'}P_{\bar{\boldsymbol{B}}}\\
&-\sum_{\boldsymbol{B},\bar{\boldsymbol{B}},\boldsymbol{B}'\in\mathbb{K}_{-1}} q^{n_2(\boldsymbol{B}'_1+\bar{\boldsymbol{B}}_1+m_1)+\boldsymbol{B}_2m_2+\boldsymbol{B}_2(\boldsymbol{B}'_1+\bar{\boldsymbol{B}}_1+m_1)+n_1\boldsymbol{B}'_1+\bar{\boldsymbol{B}}_2m_1+\bar{\boldsymbol{B}}_2\boldsymbol{B}'_1}\\
&\cdot Q_{\boldsymbol{B}+\boldsymbol{B}'+\bar{\boldsymbol{B}}+(3m_1+1,3n_1+1)+(3m_2+1,3n_2+1)}P_{\boldsymbol{B}}P_{\boldsymbol{B}'}P_{\bar{\boldsymbol{B}}} \qquad \blacksquare
\end{aligned}
$$

$$
\begin{aligned}
&-\sum_{\boldsymbol{B},\bar{\boldsymbol{B}},\boldsymbol{B}'\in\mathbb{K}_{-1}} q^{n_2\boldsymbol{B}'_1+(\boldsymbol{B}_2+\bar{\boldsymbol{B}}_2+n_1)m_2+(\boldsymbol{B}_2+\bar{\boldsymbol{B}}_2+n_1)\boldsymbol{B}'_1+n_1\bar{\boldsymbol{B}}_1+\boldsymbol{B}_2m_1+\boldsymbol{B}_2\bar{\boldsymbol{B}}_1}\\
&\cdot Q_{\boldsymbol{B}+\boldsymbol{B}'+\bar{\boldsymbol{B}}+(3m_1+1,3n_1+1)+(3m_2+1,3n_2+1)}P_{\boldsymbol{B}}P_{\boldsymbol{B}'}P_{\bar{\boldsymbol{B}}}\\
&+\sum_{\substack{\boldsymbol{A}\in\mathbb{K}_1\\ \boldsymbol{B},\boldsymbol{B}'\in\mathbb{K}_{-1}}} q^{n_1(\boldsymbol{B}_1+\boldsymbol{B}'_1+m_2)+\boldsymbol{A}_2m_1+\boldsymbol{A}_2(\boldsymbol{B}_1+\boldsymbol{B}'_1+m_2)+n_2\boldsymbol{B}'_1+\boldsymbol{B}_2m_2+\boldsymbol{B}_2\boldsymbol{B}'_1}\\
&\cdot Q_{\boldsymbol{A}+\boldsymbol{B}+\boldsymbol{B}'+(3m_1+1,3n_1+1)+(3m_2+1,3n_2+1)}P_{\boldsymbol{A}}P_{\boldsymbol{B}}P_{\boldsymbol{B}'}\\
&+\sum_{\substack{\boldsymbol{A}\in\mathbb{K}_1\\ \boldsymbol{B},\boldsymbol{B}'\in\mathbb{K}_{-1}}} q^{n_1\boldsymbol{B}_1+(\boldsymbol{A}_2+\boldsymbol{B}'_2+n_2)m_1+(\boldsymbol{A}_2+\boldsymbol{B}'_2+n_2)\boldsymbol{B}_1+n_2\boldsymbol{B}'_1+\boldsymbol{A}_2m_2+\boldsymbol{A}_2\boldsymbol{B}'_1}\\
&\cdot Q_{\boldsymbol{A}+\boldsymbol{B}+\boldsymbol{B}'+(3m_1+1,3n_1+1)+(3m_2+1,3n_2+1)}P_{\boldsymbol{A}}P_{\boldsymbol{B}}P_{\boldsymbol{B}'}\\
&-\sum_{\substack{\boldsymbol{A}\in\mathbb{K}_1\\ \boldsymbol{B},\boldsymbol{B}'\in\mathbb{K}_{-1}}} q^{n_2(\boldsymbol{B}_1+\boldsymbol{B}'_1+m_1)+\boldsymbol{A}_2m_2+\boldsymbol{A}_2(\boldsymbol{B}_1+\boldsymbol{B}'_1+m_1)+n_1\boldsymbol{B}'_1+\boldsymbol{B}_2m_1+\boldsymbol{B}_2\boldsymbol{B}'_1}\\
&\cdot Q_{\boldsymbol{A}+\boldsymbol{B}+\boldsymbol{B}'+(3m_1+1,3n_1+1)+(3m_2+1,3n_2+1)}P_{\boldsymbol{A}}P_{\boldsymbol{B}}P_{\boldsymbol{B}'}\\
&-\sum_{\substack{\boldsymbol{A}\in\mathbb{K}_1\\ \boldsymbol{B},\boldsymbol{B}'\in\mathbb{K}_{-1}}} q^{n_2\boldsymbol{B}_1+(\boldsymbol{A}_2+\boldsymbol{B}'_2+n_1)m_2+(\boldsymbol{A}_2+\boldsymbol{B}'_2+n_1)\boldsymbol{B}_1+n_1\boldsymbol{B}'_1+\boldsymbol{A}_2m_1+\boldsymbol{A}_2\boldsymbol{B}'_1}\\
&\cdot Q_{\boldsymbol{A}+\boldsymbol{B}+\boldsymbol{B}'+(3m_1+1,3n_1+1)+(3m_2+1,3n_2+1)}P_{\boldsymbol{A}}P_{\boldsymbol{B}}P_{\boldsymbol{B}'}=0
\end{aligned}
$$

as the first two terms, the third and the sixth, the fourth and the fifth, the seventh and the last term, are negative to each other.

$$
\begin{aligned}
&[e^{(\mu)}_{23}(m_1,n_1),e^{(\mu)}_{31}(m_2,n_2)]\\
=&[-q^{-m_1n_1}\mu P_{-(3m_1+1,3n_1+1)}-\sum_{\substack{\boldsymbol{A}\in\mathbb{K}_1\\ \boldsymbol{B}\in\mathbb{K}_{-1}}} q^{n_1\boldsymbol{B}_1+\boldsymbol{A}_2m_1+\boldsymbol{A}_2\boldsymbol{B}_1}Q_{\boldsymbol{A}+\boldsymbol{B}+(3m_1+1,3n_1+1)}P_{\boldsymbol{A}}P_{\boldsymbol{B}}\\
&-\sum_{\boldsymbol{B},\boldsymbol{B}'\in\mathbb{K}_{-1}} q^{n_1\boldsymbol{B}'_1+\boldsymbol{B}_2m_1+\boldsymbol{B}_2\boldsymbol{B}'_1}Q_{\boldsymbol{B}+\boldsymbol{B}'+(3m_1+1,3n_1+1)}P_{\boldsymbol{B}}P_{\boldsymbol{B}'},\\
&\sum_{\boldsymbol{A}\in\mathbb{K}_1} q^{\boldsymbol{A}_1n_2}Q_{(3m_2-2,3n_2-2)+\boldsymbol{A}}P_{\boldsymbol{A}}]
\end{aligned}
$$

$$\begin{aligned}
=&-q^{-m_1n_1+(-m_1-m_2)n_2}\mu P_{-(3m_1+1,3n_1+1)-(3m_2-2,3n_2-2)}\\
&-\sum_{\boldsymbol{A},\boldsymbol{A'}\in\mathbb{K}_1} q^{n_1(m_2+\boldsymbol{A}'_1)+\boldsymbol{A}_2m_1+\boldsymbol{A}_2(m_2+\boldsymbol{A}'_1)+\boldsymbol{A}'_1n_2}Q_{\boldsymbol{A}+\boldsymbol{A'}+(3m_2-2,3n_2-2)+(3m_1+1,3n_1+1)}P_{\boldsymbol{A}}P_{\boldsymbol{A'}}\\
&+\sum_{\substack{\boldsymbol{A}\in\mathbb{K}_1\\ \boldsymbol{B}\in\mathbb{K}_{-1}}} q^{(\boldsymbol{A}_1+\boldsymbol{B}_1+m_1)n_2+\boldsymbol{A}_2m_1+\boldsymbol{A}_2\boldsymbol{B}_1+n_1\boldsymbol{B}_1}Q_{\boldsymbol{A}+\boldsymbol{B}+(3m_2-2,3n_2-2)+(3m_1+1,3n_1+1)}P_{\boldsymbol{A}}P_{\boldsymbol{B}}\\
&-\sum_{\substack{\boldsymbol{A}\in\mathbb{K}_1\\ \boldsymbol{B}\in\mathbb{K}_{-1}}} q^{n_1(m_2+\boldsymbol{A}_1)+\boldsymbol{B}_2m_1+\boldsymbol{B}_2(m_2+\boldsymbol{A}_1)+\boldsymbol{A}_1n_2}Q_{\boldsymbol{A}+\boldsymbol{B}+(3m_2-2,3n_2-2)+(3m_1+1,3n_1+1)}P_{\boldsymbol{A}}P_{\boldsymbol{B}}\\
&-\sum_{\substack{\boldsymbol{A}\in\mathbb{K}_1\\ \boldsymbol{B}\in\mathbb{K}_{-1}}} q^{n_1\boldsymbol{B}_1+(\boldsymbol{A}_2+n_2)m_1+(\boldsymbol{A}_2+n_2)\boldsymbol{B}_1+\boldsymbol{A}_1n_2}Q_{\boldsymbol{A}+\boldsymbol{B}+(3m_2-2,3n_2-2)+(3m_1+1,3n_1+1)}P_{\boldsymbol{A}}P_{\boldsymbol{B}}
\end{aligned}$$

$$\blacksquare$$

(the third term and the fifth are cancelled)

$$\begin{aligned}
=&q^{n_1m_2}(-q^{-(m_1+m_2)(n_1+n_2)}\mu P_{-(3m_1+3m_2-1,3n_1+3n_2-1)}\\
&-\sum_{\boldsymbol{A},\boldsymbol{A'}\in\mathbb{K}_1} q^{(n_1+n_2)\boldsymbol{A}'_1+\boldsymbol{A}_2(m_1+m_2)+\boldsymbol{A}_2\boldsymbol{A}'_1}Q_{\boldsymbol{A}+\boldsymbol{A'}+(3m_2+3m_1-1,3n_2+3n_1-1)}P_{\boldsymbol{A}}P_{\boldsymbol{A'}}\\
&-\sum_{\substack{\boldsymbol{A}\in\mathbb{K}_1\\ \boldsymbol{B}\in\mathbb{K}_{-1}}} q^{(n_1+n_2)\boldsymbol{A}_1+\boldsymbol{B}_2(m_1+m_2)+\boldsymbol{B}_2\boldsymbol{A}_1}Q_{\boldsymbol{A}+\boldsymbol{B}+(3m_2+3m_1-1,3n_2+3n_1-1)}P_{\boldsymbol{A}}P_{\boldsymbol{B}})\\
=&q^{n_1m_2}e^{(\mu)}_{21}(m_1+m_2,n_1+n_2).
\end{aligned}$$

$$\begin{aligned}
&[e^{(\mu)}_{23}(m_1,n_1),e^{(\mu)}_{32}(m_2,n_2)]\\
=&[-\mu q^{-m_1n_1}P_{-(3m_1+1,3n_1+1)}-\sum_{\substack{\boldsymbol{A}\in\mathbb{K}_1\\ \boldsymbol{B}\in\mathbb{K}_{-1}}} q^{n_1\boldsymbol{B}_1+\boldsymbol{A}_2m_1+\boldsymbol{A}_2\boldsymbol{B}_1}Q_{(3m_1+1,3n_1+1)+\boldsymbol{A}+\boldsymbol{B}}P_{\boldsymbol{A}}P_{\boldsymbol{B}}\\
&-\sum_{\boldsymbol{B},\boldsymbol{B'}\in\mathbb{K}_{-1}} q^{n_1\boldsymbol{B}'_1+\boldsymbol{B}_2m_1+\boldsymbol{B}_2\boldsymbol{B}'_1}Q_{(3m_1+1,3n_1+1)+\boldsymbol{B}+\boldsymbol{B'}}P_{\boldsymbol{B}}P_{\boldsymbol{B'}},Q_{(3m_2-1,3n_2-1)}]\\
=&-\mu q^{-m_1n_1}\delta_{(-m_1,-n_1),(m_2,n_2)}\\
&-\sum_{\boldsymbol{A}\in\mathbb{K}_1} q^{n_1m_2+\boldsymbol{A}_2m_1+\boldsymbol{A}_2m_2}Q_{(3m_1+1,3n_1+1)+\boldsymbol{A}+(3m_2-1,3n_2-1)}P_{\boldsymbol{A}}\\
&-\sum_{\boldsymbol{B}\in\mathbb{K}_{-1}} q^{n_1m_2+\boldsymbol{B}_2m_1+\boldsymbol{B}_2m_2}Q_{(3m_1+1,3n_1+1)+\boldsymbol{B}+(3m_2-1,3n_2-1)}P_{\boldsymbol{B}}\\
&-\sum_{\boldsymbol{B'}\in\mathbb{K}_{-1}} q^{n_1\boldsymbol{B}'_1+n_2m_1+n_2\boldsymbol{B}'_1}Q_{(3m_1+1,3n_1+1)+\boldsymbol{B'}+(3m_2-1,3n_2-1)}P_{\boldsymbol{B'}}
\end{aligned}$$

$$
\begin{aligned}
&=q^{n_1m_2}(-\sum_{\boldsymbol{A}\in\mathbb{K}_1} q^{\boldsymbol{A}_2(m_1+m_2)}Q_{(3m_1+3m_2,3n_1+3n_2)+\boldsymbol{A}}P_{\boldsymbol{A}}\\
&\quad-\sum_{\boldsymbol{B}\in\mathbb{K}_{-1}} q^{\boldsymbol{B}_2(m_1+m_2)}Q_{(3m_1+3m_2,3n_1+3n_2)+\boldsymbol{B}}P_{\boldsymbol{B}}-\frac{1}{2}\mu\delta_{(m_1+m_2,n_1+n_2),(0,0)})\\
&\quad-q^{n_2m_1}(\sum_{\boldsymbol{B}\in\mathbb{K}_{-1}} q^{\boldsymbol{B}_1(n_1+n_2)}Q_{(3m_1+3m_2,3n_1+3n_2)+\boldsymbol{B}}P_{\boldsymbol{B}}+\frac{1}{2}\mu\delta_{(m_1+m_2,n_1+n_2),(0,0)})\\
&=q^{n_1m_2}e^{(\mu)}_{22}(m_1+m_2,n_1+n_2)-q^{n_2m_1}e^{(\mu)}_{33}(m_1+m_2,n_1+n_2).
\end{aligned}
$$

$$
\begin{aligned}
&[e^{(\mu)}_{23}(m_1,n_1),e^{(\mu)}_{33}(m_2,n_2)]\\
=&[-\mu q^{-m_1n_1}P_{-(3m_1+1,3n_1+1)}-\sum_{\substack{\boldsymbol{A}\in\mathbb{K}_1\\ \boldsymbol{B}\in\mathbb{K}_{-1}}} q^{n_1\boldsymbol{B}_1+\boldsymbol{A}_2m_1+\boldsymbol{A}_2\boldsymbol{B}_1}Q_{(3m_1+1,3n_1+1)+\boldsymbol{A}+\boldsymbol{B}}P_{\boldsymbol{A}}P_{\boldsymbol{B}}\\
&-\sum_{\boldsymbol{B},\boldsymbol{B}'\in\mathbb{K}_{-1}} q^{n_1\boldsymbol{B}'_1+\boldsymbol{B}_2m_1+\boldsymbol{B}_2\boldsymbol{B}'_1}Q_{(3m_1+1,3n_1+1)+\boldsymbol{B}+\boldsymbol{B}'}P_{\boldsymbol{B}}P_{\boldsymbol{B}'},\\
&\sum_{\boldsymbol{B}\in\mathbb{K}_{-1}} q^{\boldsymbol{B}_1n_2}Q_{(3m_2,3n_2)+\boldsymbol{B}}P_{\boldsymbol{B}}]\\
=&-q^{-m_1n_1+(-m_1-m-2)n_2}\mu P_{-(3m_1+1,3n_1+1)-(3m_2,3n_2)}\\
&-\sum_{\substack{\boldsymbol{A}\in\mathbb{K}_1\\ \boldsymbol{B}\in\mathbb{K}_{-1}}} q^{n_1(m_2+\boldsymbol{B}_1)+\boldsymbol{A}_2m_1+\boldsymbol{B}_2(m_2+\boldsymbol{B}_1)+\boldsymbol{B}_1n_2}Q_{\boldsymbol{A}+\boldsymbol{B}+(3m_1+1,3n_1+1)+(3m_2,3n_2)}P_{\boldsymbol{A}}P_{\boldsymbol{B}}\\
&-\sum_{\boldsymbol{B},\boldsymbol{B}'\in\mathbb{K}_{-1}} q^{n_1(m_2+\boldsymbol{B}'_1)+\boldsymbol{B}_2m_1+\boldsymbol{B}_2(m_2+\boldsymbol{B}'_1)+\boldsymbol{B}'_1n_2}Q_{\boldsymbol{B}+(3m_2,3n_2)+\boldsymbol{B}'+(3m_1+1,3n_1+1)}P_{\boldsymbol{B}}P_{\boldsymbol{B}'}\\
&-\sum_{\boldsymbol{B},\boldsymbol{B}'\in\mathbb{K}_{-1}} q^{n_1\boldsymbol{B}'_1+(n_2+\boldsymbol{B}_2)m_1+(\boldsymbol{B}_2+n_2)\boldsymbol{B}'_1+\boldsymbol{B}_1n_2}Q_{\boldsymbol{B}+(3m_2,3n_2)+\boldsymbol{B}'+(3m_1+1,3n_1+1)}P_{\boldsymbol{B}}P_{\boldsymbol{B}'}\\
&+\sum_{\boldsymbol{B},\boldsymbol{B}'\in\mathbb{K}_{-1}} q^{(\boldsymbol{B}_1+\boldsymbol{B}'_1+m_1)n_2+n_1\boldsymbol{B}'_1+\boldsymbol{B}_2m_1+\boldsymbol{B}_2\boldsymbol{B}'_1}Q_{\boldsymbol{B}+(3m_2,3n_2)+\boldsymbol{B}'+(3m_1+1,3n_1+1)}P_{\boldsymbol{B}}P_{\boldsymbol{B}'}
\end{aligned}
$$

(the last two terms cancel)

$$
\begin{aligned}
&=q^{n_1m_2}(-q^{-(m_1+m_2)(n_1+n_2)}\mu P_{-(3m_1+3m_2+1,3n_1+3n_2+1)}\\
&\quad-\sum_{\substack{\boldsymbol{A}\in\mathbb{K}_1\\ \boldsymbol{B}\in\mathbb{K}_{-1}}} q^{(n_1+n_2)\boldsymbol{B}_1+\boldsymbol{A}_2(m_1+m_2)+\boldsymbol{A}_2\boldsymbol{B}_1}Q_{\boldsymbol{A}+\boldsymbol{B}+(3m_1+3m_2+1,3n_1+3n_2+1)}P_{\boldsymbol{A}}P_{\boldsymbol{B}}\\
&\quad-\sum_{\boldsymbol{B},\boldsymbol{B}'\in\mathbb{K}_{-1}} q^{(n_1+n_2)\boldsymbol{B}'_1+\boldsymbol{B}_2(m_1+m_2)+\boldsymbol{B}_2\boldsymbol{B}'_1}Q_{\boldsymbol{B}+\boldsymbol{B}'+(3m_1+3m_2+1,3n_1+3n_2+1)}P_{\boldsymbol{B}}P_{\boldsymbol{B}'})\\
&=q^{n_1m_2}e^{(\mu)}_{23}(m_1+m_2,n_1+n_2).
\end{aligned}
$$

The following five brackets are easy.

$$\begin{aligned}
&[e_{31}^{(\mu)}(m_1,n_1),e_{31}^{(\mu)}(m_2,n_2)]=0\\
&[e_{31}^{(\mu)}(m_1,n_1),e_{32}^{(\mu)}(m_2,n_2)]=0\\
&[e_{31}^{(\mu)}(m_1,n_1),e_{33}^{(\mu)}(m_2,n_2)]=-q^{m_1n_2}e_{31}^{(\mu)}(m_1+m_2,n_1+n_2)\\
&[e_{32}^{(\mu)}(m_1,n_1),e_{32}^{(\mu)}(m_2,n_2)]=0\\
&[e_{32}^{(\mu)}(m_1,n_1),e_{33}^{(\mu)}(m_2,n_2)]=-q^{m_1n_2}e_{32}^{(\mu)}(m_1+m_2,n_1+n_2).
\end{aligned}$$

$$\begin{aligned}
&[e_{33}^{(\mu)}(m_1,n_1),e_{33}^{(\mu)}(m_2,n_2)]\\
=&[\sum_{\boldsymbol{B}\in\mathbb{K}_{-1}}q^{\boldsymbol{B}_1n_1}Q_{(3m_1,3n_1)+\boldsymbol{B}}P_{\boldsymbol{B}},\sum_{\boldsymbol{B}\in\mathbb{K}_{-1}}q^{\boldsymbol{B}_1n_2}Q_{(3m_2,3n_2)+\boldsymbol{B}}P_{\boldsymbol{B}}]\\
=&\sum_{\boldsymbol{B}\in\mathbb{K}_{-1}}q^{(\boldsymbol{B}_1+m_2)n_1+\boldsymbol{B}_1n_2}Q_{(3m_1,3n_1)+(3m_2,3n_2)+\boldsymbol{B}}P_{\boldsymbol{B}}\\
&-\sum_{\boldsymbol{B}\in\mathbb{K}_{-1}}q^{(\boldsymbol{B}_1+m_1)n_2+\boldsymbol{B}_1n_1}Q_{(3m_1,3n_1)+(3m_2,3n_2)+\boldsymbol{B}}P_{\boldsymbol{B}}\\
=&q^{n_1m_2}\sum_{\boldsymbol{B}\in\mathbb{K}_{-1}}q^{\boldsymbol{B}_1(n_1+n_2}Q_{(3m_1,3n_1)+(3m_2,3n_2)+\boldsymbol{B}}P_{\boldsymbol{B}}\\
&-q^{m_1n_2}\sum_{\boldsymbol{B}\in\mathbb{K}_{-1}}q^{\boldsymbol{B}_1(n_1+n_2}Q_{(3m_1,3n_1)+(3m_2,3n_2)+\boldsymbol{B}}P_{\boldsymbol{B}}\\
=&q^{n_1m_2}e_{33}^{(\mu)}(m_1+m_2,n_1+n_2)-q^{n_2m_1}e_{33}^{(\mu)}(m_1+m_2,n_1+n_2).
\end{aligned}$$

Next we check the brackets involving $D_1^{(\mu)}$ and $D_2^{(\mu)}$.

$$\begin{aligned}
&[D_1^{(\mu)},e_{11}^{(\mu)}(m_1,n_1)]\\
=&[\sum_{\boldsymbol{A}\in\mathbb{K}_1}\boldsymbol{A}_1Q_{\boldsymbol{A}}P_{\boldsymbol{A}}+\sum_{\boldsymbol{B}\in\mathbb{K}_{-1}}\boldsymbol{B}_1Q_{\boldsymbol{B}}P_{\boldsymbol{B}},\sum_{\boldsymbol{A}\in\mathbb{K}_1}q^{\boldsymbol{A}_1n_1}Q_{(3m_1,3n_1)+\boldsymbol{A}}P_{\boldsymbol{A}}+\frac{1}{2}\mu\delta_{(m_1,n_1),(0,0)}]\\
=&\sum_{\boldsymbol{A}\in\mathbb{K}_1}(m_1+\boldsymbol{A}_1)q^{\boldsymbol{A}_1n_1}Q_{(3m_1,3n_1)+\boldsymbol{A}}P_{\boldsymbol{A}}-\sum_{\boldsymbol{A}\in\mathbb{K}_1}\boldsymbol{A}_1q^{\boldsymbol{A}_1n_1}Q_{(3m_1,3n_1)+\boldsymbol{A}}P_{\boldsymbol{A}}\\
=&m_1(\sum_{\boldsymbol{A}\in\mathbb{K}_1}q^{\boldsymbol{A}_1n_1}Q_{(3m_1,3n_1)+\boldsymbol{A}}P_{\boldsymbol{A}}+\frac{1}{2}\mu\delta_{(m_1,n_1),(0,0)})\\
=&m_1e_{11}^{(\mu)}(m_1,n_1).
\end{aligned}$$

The following two brackets are easy.

$$
\begin{aligned}
[D_1^{(\mu)}, e_{12}^{(\mu)}(m_1,n_1)] &= m_1 e_{12}^{(\mu)}(m_1,n_1)\\
[D_1^{(\mu)}, e_{13}^{(\mu)}(m_1,n_1)] &= m_1 e_{13}^{(\mu)}(m_1,n_1).
\end{aligned}
$$

$$
\begin{aligned}
&[D_1^{(\mu)}, e_{21}^{(\mu)}(m_1,n_1)]\\
=&[\sum_{\boldsymbol{A}\in\mathbb{K}_1}\boldsymbol{A}_1 Q_{\boldsymbol{A}}P_{\boldsymbol{A}}+\sum_{\boldsymbol{B}\in\mathbb{K}_{-1}}\boldsymbol{B}_1 Q_{\boldsymbol{B}}P_{\boldsymbol{B}}, -q^{-m_1n_1}\mu P_{-(3m_1-1,3n_1-1)}\\
&-\sum_{\boldsymbol{A},\boldsymbol{A}'\in\mathbb{K}_1} q^{n_1\boldsymbol{A}'_1+\boldsymbol{A}_2m_1+\boldsymbol{A}_2\boldsymbol{A}'_1}Q_{\boldsymbol{A}+\boldsymbol{A}'+(3m_1-1,3n_1-1)}P_{\boldsymbol{A}}P_{\boldsymbol{A}'}\\
&-\sum_{\substack{\boldsymbol{A}\in\mathbb{K}_1\\ \boldsymbol{B}\in\mathbb{K}_{-1}}} q^{n_1\boldsymbol{A}_1+\boldsymbol{B}_2m_1+\boldsymbol{B}_2\boldsymbol{A}_1}Q_{\boldsymbol{A}+\boldsymbol{B}+(3m_1-1,3n_1-1)}P_{\boldsymbol{A}}P_{\boldsymbol{B}}]\\
=&q^{-m_1n_1}\mu(-m_1)P_{-(3m_1-1,3n_1-1)}\\
&-\sum_{\boldsymbol{A},\boldsymbol{A}'\in\mathbb{K}_1}(\boldsymbol{A}_1+\boldsymbol{A}'_1+m_1) q^{n_1\boldsymbol{A}'_1+\boldsymbol{A}_2m_1+\boldsymbol{A}_2\boldsymbol{A}'_1}Q_{\boldsymbol{A}+\boldsymbol{A}'+(3m_1-1,3n_1-1)}P_{\boldsymbol{A}}P_{\boldsymbol{A}'}\\
&+\sum_{\boldsymbol{A},\boldsymbol{A}'\in\mathbb{K}_1}\boldsymbol{A}'_1 q^{n_1\boldsymbol{A}'_1+\boldsymbol{A}_2m_1+\boldsymbol{A}_2\boldsymbol{A}'_1}Q_{\boldsymbol{A}+\boldsymbol{A}'+(3m_1-1,3n_1-1)}P_{\boldsymbol{A}}P_{\boldsymbol{A}'}\\
&+\sum_{\boldsymbol{A},\boldsymbol{A}'\in\mathbb{K}_1}\boldsymbol{A}_1 q^{n_1\boldsymbol{A}'_1+\boldsymbol{A}_2m_1+\boldsymbol{A}_2\boldsymbol{A}'_1}Q_{\boldsymbol{A}+\boldsymbol{A}'+(3m_1-1,3n_1-1)}P_{\boldsymbol{A}}P_{\boldsymbol{A}'}\\
&-\sum_{\substack{\boldsymbol{A}\in\mathbb{K}_1\\ \boldsymbol{B}\in\mathbb{K}_{-1}}}(\boldsymbol{A}_1+\boldsymbol{B}_1+m_1) q^{n_1\boldsymbol{A}_1+\boldsymbol{B}_2m_1+\boldsymbol{B}_2\boldsymbol{A}_1}Q_{\boldsymbol{A}+\boldsymbol{B}+(3m_1-1,3n_1-1)}P_{\boldsymbol{A}}P_{\boldsymbol{B}}\\
&+\sum_{\substack{\boldsymbol{A}\in\mathbb{K}_1\\ \boldsymbol{B}\in\mathbb{K}_{-1}}}\boldsymbol{B}_1 q^{n_1\boldsymbol{A}_1+\boldsymbol{B}_2m_1+\boldsymbol{B}_2\boldsymbol{A}_1}Q_{\boldsymbol{A}+\boldsymbol{B}+(3m_1-1,3n_1-1)}P_{\boldsymbol{A}}P_{\boldsymbol{B}}\\
&+\sum_{\substack{\boldsymbol{A}\in\mathbb{K}_1\\ \boldsymbol{B}\in\mathbb{K}_{-1}}}\boldsymbol{A}_1 q^{n_1\boldsymbol{A}_1+\boldsymbol{B}_2m_1+\boldsymbol{B}_2\boldsymbol{A}_1}Q_{\boldsymbol{A}+\boldsymbol{B}+(3m_1-1,3n_1-1)}P_{\boldsymbol{A}}P_{\boldsymbol{B}}\\
=&m_1(-q^{-m_1n_1}\mu P_{-(3m_1-1,3n_1-1)}\\
&-\sum_{\boldsymbol{A},\boldsymbol{A}'\in\mathbb{K}_1} q^{n_1\boldsymbol{A}'_1+\boldsymbol{A}_2m_1+\boldsymbol{A}_2\boldsymbol{A}'_1}Q_{\boldsymbol{A}+\boldsymbol{A}'+(3m_1-1,3n_1-1)}P_{\boldsymbol{A}}P_{\boldsymbol{A}'}\\
&-\sum_{\substack{\boldsymbol{A}\in\mathbb{K}_1\\ \boldsymbol{B}\in\mathbb{K}_{-1}}} q^{n_1\boldsymbol{A}_1+\boldsymbol{B}_2m_1+\boldsymbol{B}_2\boldsymbol{A}_1}Q_{\boldsymbol{A}+\boldsymbol{B}+(3m_1-1,3n_1-1)}P_{\boldsymbol{A}}P_{\boldsymbol{B}})\\
=&m_1 e_{21}^{(\mu)}(m_1,n_1).
\end{aligned}
$$

$$
\begin{aligned}
&[D_1^{(\mu)}, e_{22}^{(\mu)}(m_1,n_1)]\\
=&[\sum_{\boldsymbol{A}\in\mathbb{K}_1}\boldsymbol{A}_1Q_{\boldsymbol{A}}P_{\boldsymbol{A}}+\sum_{\boldsymbol{B}\in\mathbb{K}_{-1}}\boldsymbol{B}_1Q_{\boldsymbol{B}}P_{\boldsymbol{B}},-\sum_{\boldsymbol{A}\in\mathbb{K}_1}q^{\boldsymbol{A}_2m_1}Q_{(3m_1,3n_1)+\boldsymbol{A}}P_{\boldsymbol{A}}\\
&-\sum_{\boldsymbol{B}\in\mathbb{K}_{-1}}q^{\boldsymbol{B}_2m_1}Q_{(3m_1,3n_1)+\boldsymbol{B}}P_{\boldsymbol{B}}-\frac{1}{2}\mu\delta_{(m_1,n_1),(0,0)}]\\
=&-\sum_{\boldsymbol{A}\in\mathbb{K}_1}(\boldsymbol{A}_1+m_1)q^{\boldsymbol{A}_2m_1}Q_{(3m_1,3n_1)+\boldsymbol{A}}P_{\boldsymbol{A}}+\sum_{\boldsymbol{A}\in\mathbb{K}_1}\boldsymbol{A}_1q^{\boldsymbol{A}_2m_1}Q_{(3m_1,3n_1)+\boldsymbol{A}}P_{\boldsymbol{A}}\\
&-\sum_{\boldsymbol{B}\in\mathbb{K}_{-1}}(m_1+\boldsymbol{B}_1)q^{\boldsymbol{B}_2m_1}Q_{(3m_1,3n_1)+\boldsymbol{B}}P_{\boldsymbol{B}}+\sum_{\boldsymbol{B}\in\mathbb{K}_{-1}}\boldsymbol{B}_1q^{\boldsymbol{B}_2m_1}Q_{(3m_1,3n_1)+\boldsymbol{B}}P_{\boldsymbol{B}}\\
=&m_1(-\sum_{\boldsymbol{A}\in\mathbb{K}_1}q^{\boldsymbol{A}_2m_1}Q_{(3m_1,3n_1)+\boldsymbol{A}}P_{\boldsymbol{A}}-\sum_{\boldsymbol{B}\in\mathbb{K}_{-1}}q^{\boldsymbol{B}_2m_1}Q_{(3m_1,3n_1)+\boldsymbol{B}}P_{\boldsymbol{B}}-\frac{1}{2}\mu\delta_{(m_1,n_1),(0,0)})\\
=&m_1e_{22}^{(\mu)}(m_1,n_1).
\end{aligned}
$$

$$
\begin{aligned}
&[D_1^{(\mu)}, e_{23}^{(\mu)}(m_1,n_1)]\\
=&[\sum_{\boldsymbol{A}\in\mathbb{K}_1}\boldsymbol{A}_1Q_{\boldsymbol{A}}P_{\boldsymbol{A}}+\sum_{\boldsymbol{B}\in\mathbb{K}_{-1}}\boldsymbol{B}_1Q_{\boldsymbol{B}}P_{\boldsymbol{B}},-q^{-m_1n_1}\mu P_{-(3m_1+1,3n_1+1)}\\
&-\sum_{\substack{\boldsymbol{A}\in\mathbb{K}_1\\ \boldsymbol{B}\in\mathbb{K}_{-1}}}q^{n_1\boldsymbol{B}_1+\boldsymbol{A}_2m_1+\boldsymbol{A}_2\boldsymbol{B}_1}Q_{\boldsymbol{A}+\boldsymbol{B}+(3m_1+1,3n_1+1)}P_{\boldsymbol{A}}P_{\boldsymbol{B}}\\
&-\sum_{\boldsymbol{B},\boldsymbol{B}'\in\mathbb{K}_{-1}}q^{n_1\boldsymbol{B}'_1+\boldsymbol{B}_2m_1+\boldsymbol{B}_2\boldsymbol{B}'_1}Q_{\boldsymbol{B}+\boldsymbol{B}'+(3m_1+1,3n_1+1)}P_{\boldsymbol{B}}P_{\boldsymbol{B}'}]\\
=&q^{-m_1n_1}\mu(-m_1)P_{-(3m_1+1,3n_1+1)}\\
&-\sum_{\substack{\boldsymbol{A}\in\mathbb{K}_1\\ \boldsymbol{B}\in\mathbb{K}_{-1}}}(\boldsymbol{A}_1+\boldsymbol{B}_1+m_1)q^{n_1\boldsymbol{B}_1+\boldsymbol{A}_2m_1+\boldsymbol{A}_2\boldsymbol{B}_1}Q_{\boldsymbol{A}+\boldsymbol{B}+(3m_1+1,3n_1+1)}P_{\boldsymbol{A}}P_{\boldsymbol{B}}\\
&+\sum_{\substack{\boldsymbol{A}\in\mathbb{K}_1\\ \boldsymbol{B}\in\mathbb{K}_{-1}}}\boldsymbol{A}_1q^{n_1\boldsymbol{B}_1+\boldsymbol{A}_2m_1+\boldsymbol{A}_2\boldsymbol{B}_1}Q_{\boldsymbol{A}+\boldsymbol{B}+(3m_1+1,3n_1+1)}P_{\boldsymbol{A}}P_{\boldsymbol{B}}\\
&+\sum_{\substack{\boldsymbol{A}\in\mathbb{K}_1\\ \boldsymbol{B}\in\mathbb{K}_{-1}}}\boldsymbol{B}_1q^{n_1\boldsymbol{B}_1+\boldsymbol{A}_2m_1+\boldsymbol{A}_2\boldsymbol{B}_1}Q_{\boldsymbol{A}+\boldsymbol{B}+(3m_1+1,3n_1+1)}P_{\boldsymbol{A}}P_{\boldsymbol{B}}\\
&-\sum_{\boldsymbol{B},\boldsymbol{B}'\in\mathbb{K}_{-1}}(\boldsymbol{B}_1+\boldsymbol{B}'_1+m_1)q^{n_1\boldsymbol{B}'_1+\boldsymbol{B}_2m_1+\boldsymbol{B}_2\boldsymbol{B}'_1}Q_{\boldsymbol{B}+\boldsymbol{B}'+(3m_1+1,3n_1+1)}P_{\boldsymbol{B}}P_{\boldsymbol{B}'}
\end{aligned}
$$

$$
\begin{aligned}
&+\sum_{\boldsymbol{B},\boldsymbol{B}'\in\mathbb{K}_{-1}} \boldsymbol{B}_1 q^{n_1\boldsymbol{B}'_1+\boldsymbol{B}_2 m_1+\boldsymbol{B}_2\boldsymbol{B}'_1} Q_{\boldsymbol{B}+\boldsymbol{B}'+(3m_1+1,3n_1+1)} P_{\boldsymbol{B}} P_{\boldsymbol{B}'} \\
&+\sum_{\boldsymbol{B},\boldsymbol{B}'\in\mathbb{K}_{-1}} \boldsymbol{B}'_1 q^{n_1\boldsymbol{B}'_1+\boldsymbol{B}_2 m_1+\boldsymbol{B}_2\boldsymbol{B}'_1} Q_{\boldsymbol{B}+\boldsymbol{B}'+(3m_1+1,3n_1+1)} P_{\boldsymbol{B}} P_{\boldsymbol{B}'} \\
=&m_1(-q^{-m_1n_1}\mu P_{-(3m_1+1,3n_1+1)} \\
&-\sum_{\substack{\boldsymbol{A}\in\mathbb{K}_1\\ \boldsymbol{B}\in\mathbb{K}_{-1}}} q^{n_1\boldsymbol{B}_1+\boldsymbol{A}_2 m_1+\boldsymbol{A}_2\boldsymbol{B}_1} Q_{\boldsymbol{A}+\boldsymbol{B}+(3m_1+1,3n_1+1)} P_{\boldsymbol{A}} P_{\boldsymbol{B}} \\
&-\sum_{\boldsymbol{B},\boldsymbol{B}'\in\mathbb{K}_{-1}} q^{n_1\boldsymbol{B}'_1+\boldsymbol{B}_2 m_1+\boldsymbol{B}_2\boldsymbol{B}'_1} Q_{\boldsymbol{B}+\boldsymbol{B}'+(3m_1+1,3n_1+1)} P_{\boldsymbol{B}} P_{\boldsymbol{B}'}) \\
=&m_1 e_{23}^{(\mu)}(m_1,n_1).
\end{aligned}
$$

The following two brackets are easy.

$$
\begin{aligned}
[D_1^{(\mu)}, e_{31}^{(\mu)}(m_1,n_1)] &= m_1 e_{31}^{(\mu)}(m_1,n_1) \\
[D_1^{(\mu)}, e_{32}^{(\mu)}(m_1,n_1)] &= m_1 e_{32}^{(\mu)}(m_1,n_1).
\end{aligned}
$$

$$
\begin{aligned}
&[D_1^{(\mu)}, e_{33}^{(\mu)}(m_1,n_1)] \\
=&[\sum_{\boldsymbol{A}\in\mathbb{K}_1} \boldsymbol{A}_1 Q_{\boldsymbol{A}} P_{\boldsymbol{A}} + \sum_{\boldsymbol{B}\in\mathbb{K}_{-1}} \boldsymbol{B}_1 Q_{\boldsymbol{B}} P_{\boldsymbol{B}}, \sum_{\boldsymbol{B}\in\mathbb{K}_{-1}} q^{\boldsymbol{B}_1 n_1} Q_{(3m_1,3n_1)+\boldsymbol{B}} P_{\boldsymbol{B}} + \frac{1}{2}\mu\delta_{(m_1,n_1),(0,0)}] \\
=&\sum_{\boldsymbol{B}\in\mathbb{K}_{-1}} (m_1+\boldsymbol{B}_1) q^{\boldsymbol{B}_1 n_1} Q_{(3m_1,3n_1)+\boldsymbol{B}} P_{\boldsymbol{B}} - \sum_{\boldsymbol{B}\in\mathbb{K}_{-1}} \boldsymbol{B}_1 q^{\boldsymbol{B}_1 n_1} Q_{(3m_1,3n_1)+\boldsymbol{B}} P_{\boldsymbol{B}} \\
=&m_1(\sum_{\boldsymbol{B}\in\mathbb{K}_{-1}} q^{\boldsymbol{B}_1 n_1} Q_{(3m_1,3n_1)+\boldsymbol{B}} P_{\boldsymbol{B}} + \frac{1}{2}\mu\delta_{(m_1,n_1),(0,0)}) \\
=&m_1 e_{33}^{(\mu)}(m_1,n_1)
\end{aligned}
$$

Similarly, we can get

$$
[D_2^{(\mu)}, e_{ij}^{(\mu)}(m_1,n_1)] = n_1 e_{ij}^{(\mu)}(m_1,n_1)
$$

for $1 \le i,j \le 3$. Finally,

$$
\begin{aligned}
&[D_1^{(\mu)}, D_2^{(\mu)}] \\
=&[\sum_{\boldsymbol{A}\in\mathbb{K}_1} \boldsymbol{A}_1 Q_{\boldsymbol{A}} P_{\boldsymbol{A}} + \sum_{\boldsymbol{B}\in\mathbb{K}_{-1}} \boldsymbol{B}_1 Q_{\boldsymbol{B}} P_{\boldsymbol{B}}, \sum_{\boldsymbol{A}\in\mathbb{K}_1} \boldsymbol{A}_2 Q_{\boldsymbol{A}} P_{\boldsymbol{A}} + \sum_{\boldsymbol{B}\in\mathbb{K}_{-1}} \boldsymbol{B}_2 Q_{\boldsymbol{B}} P_{\boldsymbol{B}}] \\
=&0
\end{aligned}
$$

Hence $\pi : \widetilde{\mathfrak{gl}_3(\mathbb{C}_q)} \to End(V)$ is a Lie algebra homomorphism. $\square$

§2. Hermitian form for $\widetilde{\mathfrak{gl}_3(\mathbb{C}_q)}$-module.

From now on we need to assume that $|q| = 1$.

Define $\omega : \widetilde{\mathfrak{gl}_3(\mathbb{C}_q)} \mapsto \widetilde{\mathfrak{gl}_3(\mathbb{C}_q)}$ a $\mathbb{R}$-linear map as the following:

$$\omega(\lambda x) = \bar{\lambda}\omega(x), \forall \lambda \in \mathbb{C}, x \in \widetilde{\mathfrak{gl}_3(\mathbb{C}_q)} \tag{2.1}$$

$$\omega(E_{ij}(a)) = (-1)^{i+j} E_{ji}(\overline{a}), a \in \mathbb{C}_q \tag{2.2}$$

$$\omega(d_s) = d_s,\, \omega(d_t) = d_t,\, \omega(c_s) = c_s,\, \omega(c_t) = c_t \tag{2.3}$$

where $\mathbb{R}-$linear function $^{-} : \mathbb{C}_q \to \mathbb{C}_q$ is defined as $\overline{\lambda s^m t^n} = \bar{\lambda} t^{-n} s^{-m} = \bar{\lambda} q^{mn} s^{-m} t^{-n}$, and $\bar{\lambda}$ is the complex conjugate, for any $\lambda \in \mathbb{C}$, and $m, n \in \mathbb{Z}$.

Following from [Lemma 3.4, GZ], we have

Lemma 2.4. *ω is an anti-linear anti-involution of $\widetilde{\mathfrak{gl}_3(\mathbb{C}_q)}$.*

We simply write $\pi(E_{ij}(r)).v$ as $E_{ij}(r).v$, for any $v \in V, r \in \mathbb{C}_q$.

In [GZ], we define a hermitian form on the basis consisting of monomials and then use another basis consisting of iterated module actions on the "highest weight vector " 1 to determine the condition for the form being positive definite. Here we will use the second basis directly to define the hermitian form which is much simpler.

Lemma 2.5. $E_{12}(\alpha_1)E_{12}(\alpha_2)...E_{12}(\alpha_k)E_{32}(\beta_1)E_{32}(\beta_2)...E_{32}(\beta_l).1$ *(We shall call it in level (k, l) in W), here $k, l \in \mathbb{Z}_+ \bigcup\{0\}$, $\alpha_i = s^{m_i}t^{n_i}, i = 1...k, \beta_j = s^{u_j}t^{v_j}$, $j = 1...l, m_i, n_i, u_j, v_j \in \mathbb{Z}$ forms a basis for V.*

Proof. Since

$$f_{\boldsymbol{A},\boldsymbol{B}} = \prod_{(m,n)\in\mathbb{Z}^2} x_{(3m+1,3n+1)}^{\boldsymbol{A}_{(m,n)}} \cdot \prod_{(m',n')\in\mathbb{Z}^2} x_{(3m'-1,3n'-1)}^{\boldsymbol{B}_{(m',n')}}$$

$\boldsymbol{A}_{(m,n)}, \boldsymbol{B}_{(m',n')} \in \mathbb{Z}_+ \cup \{0\}$, where only finitely many $\boldsymbol{A}_{(m,n)}, \boldsymbol{B}_{(m',n')}$ are nonzero, form a basis for V.

Let $g_{\boldsymbol{A}} = \prod_{(m,n)\in\mathbb{Z}^2} x_{(3m+1,3n+1)}^{\boldsymbol{A}_{(m,n)}}$, and $h_{\boldsymbol{B}} = \prod_{(m',n')\in\mathbb{Z}^2} x_{(3m'-1,3n'-1)}^{\boldsymbol{B}_{(m',n')}}$. It is similar to [Lemma 4.2, GZ], $g_{\boldsymbol{A}}$ can be written as a linear combination of $E_{12}(\alpha_1)...E_{12}(\alpha_k).1$, for $k \le \sum_{(m,n)} \boldsymbol{A}_{(m,n)}$ and $h_{\boldsymbol{B}}$ can be written as a linear combination of $E_{32}(\beta_1)...E_{32}(\beta_l).1$, for $l \le \sum_{(m',n')} \boldsymbol{B}_{(m',n')}$.

Since $E_{12}(\alpha)E_{32}(\beta).u = E_{32}(\beta)E_{12}(\alpha).u$ for any $u \in V$, $f_{(\boldsymbol{A},\boldsymbol{B})}$ can be written as a linear combination of $E_{12}(\alpha_1)...E_{12}(\alpha_k)E_{32}(\beta_1)...E_{32}(\beta_l).1$. Hence

$$E_{12}(\alpha_1)...E_{12}(\alpha_k)E_{32}(\beta_1)...E_{32}(\beta_l).1$$

form a basis for V. $\square$

Let

$$\mathcal{B} = \{E_{12}(\alpha_1)...E_{12}(\alpha_k)E_{32}(\beta_1)...E_{32}(\beta_l).1| \text{ for all } k, l \in \mathbb{N}, \alpha_i, \beta_j \in \mathbb{C}_q\} \tag{2.6}$$

be the basis for V.

Lemma 2.7. *For any* $v \in V$, $lev(v) = lev(E_{ii}(a).v)$, $i = 1, 2, 3$; $lev(E_{12}(a)(v)) = lev(v)+(1, 0)$; $lev(E_{32}(a).v) = lev(v)+(0, 1)$; $lev(E_{21}(a).v) = lev(v)-(1, 0)$ *or* $E_{21}(a).v = 0$ *if* $lev(v)-(1, 0) \notin \mathbb{Z}_+^2$; $lev(E_{23}(a).v) = lev(v)-(0, 1)$ *or* $E_{23}(a).v = 0$ *if* $lev(v)-(0, 1) \notin \mathbb{Z}_+^2$, *for any* $0 \neq a \in \mathbb{C}_q$.

Proof. We only check those v in the basis $\mathcal{B}$.

$$\begin{aligned}
&E_{22}(a)E_{12}(\alpha_1)E_{12}(\alpha_2)...E_{12}(\alpha_k)E_{32}(\beta_1)E_{32}(\beta_2)...E_{32}(\beta_l).1\\
=&E_{12}(\alpha_1)E_{22}(a)E_{12}(\alpha_2)...E_{12}(\alpha_k)E_{32}(\beta_1)E_{32}(\beta_2)...E_{32}(\beta_l).1\\
&- E_{12}(\alpha_1 a)E_{12}(\alpha_2)...E_{12}(\alpha_k)E_{32}(\beta_1)E_{32}(\beta_2)...E_{32}(\beta_l).1\\
=&E_{12}(\alpha_1)E_{12}(\alpha_2)...E_{12}(\alpha_k)E_{32}(\beta_1)E_{32}(\beta_2)...E_{32}(\beta_l).(\frac{1}{2}\mu)\kappa(a).1\\
&- \sum_{i=1}^{k} E_{12}(\alpha_1)...E_{12}(\alpha_i a)...E_{12}(\alpha_k)E_{32}(\beta_1)E_{32}(\beta_2)...E_{32}(\beta_l).1\\
&+ \sum_{i=1}^{l} E_{12}(\alpha_1)E_{12}(\alpha_2)...E_{12}(\alpha_k)E_{32}(\beta_1)..E_{32}(\beta_i a)...E_{32}(\beta_l).1,
\end{aligned}$$

so $lev(v) = lev(E_{22}(a).v)$. It is similar for $E_{11}(a), E_{33}(a)$.

$lev(E_{12}(a)(v)) = lev(v) + (1, 0)$ and $lev(E_{32}(a).v) = lev(v) + (0, 1)$ are the definition of level.

For $E_{21}(a).v$, we prove by induction on the level of v:

$E_{21}(a).v = 0$ if $lev(v) = (0, n)$, $n \in \mathbb{Z}_+ \cup \{0\}$:

If $n = 0$, it is obvious that $E_{21}(a).1 = 0$.

Suppose it is true for n, then

$$\begin{aligned}
&E_{21}(a)E_{32}(\beta_1)E_{32}(\beta_2)...E_{32}(\beta_{n+1}).1\\
=&E_{32}(\beta_1)E_{21}(a)E_{32}(\beta_2)...E_{32}(\beta_{n+1}).1 - E_{31}(\beta_1 a)E_{32}(\beta_2)...E_{32}(\beta_{n+1}).1\\
=&- E_{32}(\beta_2)...E_{32}(\beta_{n+1})E_{31}(\beta_1 a).1\\
=&0
\end{aligned}$$

by induction.

Suppose $lev(E_{21}(a).v) = lev(v) - (1,0)$ or $E_{21}(a).v = 0$ is true for the $lev(v) = (m-1,n)$, then for $v = E_{12}(b)v'$ with $lev(v') = (m-1,n)$, and $0 \neq b \in \mathbb{C}_q$

$$E_{21}(a).E_{12}(b)v' = E_{12}(b)E_{21}(a)v' + E_{22}(ab).v' - E_{11}(ba).v'.$$

Since $lev(E_{21}(a)v') = (m-2,n)$, $lev(E_{21}(a).E_{12}(b)v') = (m-1,n)$ or $E_{21}(a).E_{12}(b)v' = 0$. It is similar for $E_{23}(a)$. $\square$

We easily define a contravariant (w.r.t. π, ω) hermitian form on V by defining on the basis $\mathcal{B}$.

Assume that μ is a real number, define the conjugate bilinear form on the elements in $\mathcal{B}$ by induction on the level:

$$(1,1) = 1, (1,f) = 0 \text{ if } lev(f) \neq (0,0) \tag{2.8}$$

Suppose for any $v \in \mathcal{B}$, (u,v) is defined for any u such that $lev(u) = (k',l')$, with $k'+l' = r-1$, if $lev(u) = (k,l)$, with $k+l = r$, then there exists a u' such that $lev(u') = (k-1,l)$, or $lev(u') = (k,l-1)$, and some $a \in \mathbb{C}_q$, such that $u = E_{12}(a)u'$ or $u = E_{32}(a)u'$. Define

$$(E_{12}(a)u',v) = (u',\omega(E_{12}(a))v), \tag{2.9}$$

$$(E_{32}(a)u',v) = (u',\omega(E_{32}(a))v). \tag{2.10}$$

Theorem 2.11. *The conjugate bilinear form defined above is a hermitian form on V.*

Proof. We have to check that $(E_{ij}(a)u,v) = (u,\omega(E_{ij}(a))v)$, for $1 \leq i,j \leq 3, a \in \mathbb{C}_q$, and $(D_i.u,v) = (u,\omega(D_i)v)$ for $i = 1,2$:

By the definition,

$$\begin{aligned}(E_{12}(a)u,v) &= (u,\omega(E_{12}(a))v),\\ (E_{32}(a)u,v) &= (u,\omega(E_{32}(a))v),\end{aligned}$$

and so

$$\begin{aligned}(E_{13}(a)u,v) =&([E_{12}(1),E_{23}(a)]u,v)\\ =&(E_{12}(1)E_{23}(a)u,v) - (E_{23}(a)E_{12}(1)u,v)\\ =&(u,\omega(E_{23}(a))\omega(E_{12}(1))v) - (u,\omega(E_{12}(1))\omega(E_{23}(a))v)\\ =&(u,-\omega([E_{23}(a),E_{12}(1)])v) = (u,\omega(E_{13}(a))v).\end{aligned}$$

Using induction on the $lev(u)$ to prove $(E_{11}(a).u,v) = (u,\omega(E_{11}(a)).v)$:

For any $v \in \mathcal{B}$,

$$(E_{11}(a)1, v) = \frac{1}{2}\mu\kappa(a)(1, v) = \frac{1}{2}\mu\kappa(a)\delta_{1,v}.$$

Since $lev(E_{11}(a).v) = lev(v)$ for any $v \in \mathcal{B}$,

$$(1, \omega(E_{11}(a)).v) = (1, E_{11}(\bar{a}).v) = \frac{1}{2}\mu\kappa(\bar{a})\delta_{1,v}.$$

Hence

$$(E_{11}(a)1, v) = (1, \omega(E_{11}(a)).v).$$

Suppose $(E_{11}(a)u, v) = (u, \omega(E_{11}(a)).v)$ holds true for any $lev(u) = (l, k)$ with $l + k = r - 1$. For $lev(u) = (l, k)$ with $l + k = r$, then $u = E_{32}(b).u'$, with $lev(u') = (l, k - 1)$,

$$\begin{aligned}
(E_{11}(a)E_{32}(b).u', v) =&(E_{32}(b)E_{11}(a).u', v)\\
=&(E_{11}(a).u', \omega(E_{32}(b)).v)\\
=&(u', \omega(E_{11}(a))\omega(E_{32}(b)).v)\\
=&(u', \omega(E_{32}(b))\omega(E_{11}(a)).v)\\
=&(E_{32}(b)u', \omega(E_{11}(a)).v)\\
=&(u, \omega(E_{11}(a)).v),
\end{aligned}$$

or $u = E_{12}(b).u'$ with $lev(u') = (l - 1, k)$,

$$\begin{aligned}
(E_{11}(a)E_{12}(b).u', v) =&(E_{12}(b)E_{11}(a).u', v) + ([E_{11}(a), E_{12}(b)].u', v)\\
=&(E_{11}(a).u', \omega(E_{12}(b)).v) + (u', \omega([E_{11}(a), E_{12}(b)]).v)\\
=&(u', \omega(E_{11}(a))\omega(E_{12}(b)).v) - (u', [\omega(E_{11}(a)), \omega(E_{12}(b))].v)\\
=&(u', \omega(E_{12}(b))\omega(E_{11}(a)).v)\\
=&(E_{12}(b)u', \omega(E_{11}(a)).v)\\
=&(u, \omega(E_{11}(a)).v).
\end{aligned}$$

Thus $(E_{11}(a).u, v) = (u, \omega(E_{11}(a)).v)$; and

$$\begin{aligned}
(E_{22}(a).u, v) =&([E_{21}(a), E_{12}(1)].u, v) + (E_{11}(a)u, v)\\
=&(E_{21}(a)E_{12}(1).u, v) - (E_{12}(1)E_{21}(a).u, v) + (E_{11}(a)u, v)\\
=&(u, \omega(E_{12}(1))\omega(E_{21}(a)).v) - (u, \omega(E_{21}(a))\omega(E_{12}(1)).v) + (u, \omega(E_{11}(a)).v)\\
=&(u, \omega([E_{21}(a), E_{12}(1)].v)) + (u, \omega(E_{11}(a)).v)\\
=&(u, \omega(E_{22}(a)).v).
\end{aligned}$$

It is similar for $(E_{33}(a).u, v) = (u, \omega(E_{33}(a)).v)$.

For D_1, D_2, we also prove by induction on the level of u:

It is obvious that $(D_1.1, v) = 0$, for any $v \in \mathcal{B}$, so $(D_1.1, 1) = (1, D_1.1) = 0$, and suppose $(1, D_1.v) = 0$ is true for those $lev(v) = (k', l')$, with $k' + l' = r > 0$, then $(1, D_1 E_{12}(s^m t^n).v) = (1, E_{12} D_1.v) + (1, m.v) = 0$, and $(1, D_1 E_{32}(s^m t^n).v) = (1, E_{32} D_1.v) + (1, m.v) = 0$. So $(D_1.1, v) = (1, D_1.v)$.

Suppose for any $v \in \mathcal{B}$, $(D_1.u, v) = (u, D_1.v)$ is true for all $lev(u) = (k', l')$ such that $k' + l' = r$, then

$$\begin{aligned}(D_1.E_{12}(s^m t^n).u, v) &= (E_{12}(s^m t^n) D_1.u, v) + (m.u, v)\\ &= (D_1.u, \omega(E_{12}(s^m t^n))v) + (u, m.v)\\ &= (u, D_1 \omega(E_{12}(s^m t^n))v) + (u, m.v)\\ &= (u, \omega(E_{12}(s^m t^n)) D_1.v)\\ &= (E_{12}(s^m t^n)u, D_1.v).\end{aligned}$$

It is similar for $(D_1.E_{32}(s^m t^n).u, v) = (E_{32}(s^m t^n)u, D_1.v)$.

Hence $(D_1.u, v) = (u, D_1.v)$, and so is $(D_2.u, v) = (u, D_2.v)$. Note that $\omega(D_i) = D_i, i = 1, 2$. $\square$

§3. Conditions for unitarity.

In this section we will determine when the hermitian form given last section is positive definite.

Let $i \in \mathbb{N}$, $\gamma = (\gamma_1, ..., \gamma_s)$ be the $s-partition$ of i. We denote $Par_s(i)$ be the set of all $s-partition$ of i.

Let $\gamma \in Par_s(N)$, we say that $\pi_1' \times \pi_2' \in S_N \times S_N$ is equivalent to $\pi_1 \times \pi_2 \in S_N \times S_N$, where S_N is the permutation group of N letters, if for all $z_1, ...z_N, w_1, ..., w_N \in \mathbb{C}_q$,

$$\kappa(z_{\pi_1'(1)} w_{\pi_2'(1)} ... z_{\pi_1'(\gamma_1)} w_{\pi_2'(\gamma_1)}) ... \kappa(z_{\pi_1'(\gamma_1 + ... \gamma_{s-1}+1)} w_{\pi_2'(\gamma_1 + ... + \gamma_{s-1}+1)} ... z_{\pi_1'(N)} w_{\pi_2'(N)})$$

can be obtained from the analogous expression for $\pi_1 \times \pi_2$ only by rotating the variables. (e.g. $\kappa(z_1 w_1 z_2 w_2 z_3 w_3) = \kappa(z_3 w_3 z_1 w_1 z_2 w_2)$).

The following lemma is due to [JK2].

Lemma 3.1. *Let* $z_1, z_2, ..z_N, w_1, w_2, ..w_N \in \mathbb{C}_q[s^{\pm 1}, t^{\pm 1}]$

(3.2)

$$\begin{aligned}&\begin{pmatrix}0 & z_1\\ 0 & 0\end{pmatrix}\begin{pmatrix}0 & z_2\\ 0 & 0\end{pmatrix}...\begin{pmatrix}0 & z_N\\ 0 & 0\end{pmatrix}\begin{pmatrix}0 & 0\\ w_1 & 0\end{pmatrix}\begin{pmatrix}0 & 0\\ w_2 & 0\end{pmatrix}...\begin{pmatrix}0 & 0\\ w_N & 0\end{pmatrix}.1\\ =&\sum_{s=1}^{N} \sum_{\gamma \in Par_s(N)} \sum_{[\pi_1 \times \pi_2] \in (S_N \times S_N)(\gamma)} (-1)^{\gamma_1 - 1}(-\mu)\kappa(z_{\pi_1(1)} w_{\pi_2(1)} ... z_{\pi_1(\gamma_1)} w_{\pi_2(\gamma_1)})\\ &.(-1)^{\gamma_2 - 1}(-\mu)\kappa(z_{\pi_1(\gamma_1+1)} w_{\pi_2(\gamma_1+1)} ... z_{\pi_1(\gamma_2)} w_{\pi_2(\gamma_2)}).\\ &...(-1)^{\gamma_s - 1}(-\mu)\kappa(z_{\pi_1(\gamma_1 + ... \gamma_{s-1}+1)} w_{\pi_2(\gamma_1 + ... + \gamma_{s-1}+1)} ... z_{\pi_1(N)} w_{\pi_2(N)}).1\end{aligned}$$

Lemma 3.3. *Let* $a_i, c_i, b_j, d_j \in \mathbb{C}_q, i = 1, ..., m, j = 1, ..., n,$ *and* $R = (a_i c_j)_{m\times m}$, $U = (b_i d_j)_{n\times n}$, *and set* $\Lambda = \begin{pmatrix} R & 0 \\ 0 & U \end{pmatrix}_{(m+n)\times(m+n)} = (\lambda_{i,j})_{(m+n)\times(m+n)}$

(3.4)

$$\begin{aligned}
&E_{21}(a_1)...E_{21}(a_m)E_{23}(b_1)...E_{23}(b_n)E_{12}(c_1)...E_{12}(c_m)E_{32}(d_1)...E_{32}(d_n).1 \\
&= \sum_{s=1}^{m+n} \sum_{\gamma\in Par_s(m+n)} \sum_{[\pi_1\times\pi_2]\in(S_{m+n}\times S_{m+n})(\gamma)} (-1)^{\gamma_1-1}(-\mu)\kappa(\lambda_{\pi_1(1),\pi_2(1)}...\lambda_{\pi_1(\gamma_1)\pi_2(\gamma_1)}) \\
&.(-1)^{\gamma_2-1}(-\mu)\kappa(\lambda_{\pi_1(\gamma_1+1),\pi_2(\gamma_1+1)}...\lambda_{\pi_1(\gamma_2),\pi_2(\gamma_2)}). \\
&...(-1)^{\gamma_s-1}(-\mu)\kappa(\lambda_{\pi_1(\gamma_1+...\gamma_{s-1}+1),\pi_2(\gamma_1+...+\gamma_{s-1}+1)}...\lambda_{\pi_1(N),\pi_2(N)}).1
\end{aligned}$$

Remark 3.5. *It is easy to see that* $\lambda_{i,j}$ *in every summand should be from different rows and different columns of* Λ. *And if the summand of (3.4) contains some* $\lambda_{i,j} = 0$, *then this summand is* 0. *Hence (3.4) in fact is the sum of those* $\lambda_{i,j}$ *from* R *and* U.

Proof. Prove by induction on n:

$n = 0$, (3.4) is just (3.2).

Assume (3.4) is true up to $n-1$,

$$\begin{aligned}
&E_{21}(a_1)...E_{21}(a_m)E_{23}(b_1)...E_{23}(b_n)E_{12}(c_1)...E_{12}(c_m)E_{32}(d_1)...E_{32}(d_n).1 \\
=&E_{21}(a_1)...E_{21}(a_m)E_{23}(b_1)...E_{23}(b_{n-1}) \\
&(E_{12}(c_1)E_{23}(b_n) - E_{13}(c_1b_n))E_{12}(c_2)...E_{12}(c_m)E_{32}(d_1)...E_{32}(d_n).1 \\
=&E_{21}(a_1)...E_{21}(a_m)E_{23}(b_1)...E_{23}(b_{n-1})E_{12}(c_1)E_{23}(b_n)E_{12}(c_2)...E_{12}(c_m)E_{32}(d_1)...E_{32}(d_n).1 \\
&- E_{21}(a_1)...E_{21}(a_m)E_{23}(b_1)...E_{23}(b_{n-1})E_{12}(c_1)...E_{12}(c_m)(E_{12}(c_1b_nd_1) \\
&+ E_{32}(d_1)E_{13}(c_1b_n))E_{32}(d_2)...E_{32}(d_n).1 \\
=&E_{21}(a_1)...E_{21}(a_m)E_{23}(b_1)...E_{23}(b_{n-1})E_{12}(c_1)E_{23}(b_n)E_{12}(c_2)...E_{12}(c_m)E_{32}(d_1)...E_{32}(d_n).1 \\
&+ \sum_{i=1}^{n} E_{21}(a_1)...E_{21}(a_m)E_{23}(b_1)...E_{23}(b_{n-1}). \\
&E_{12}(-c_1b_nd_i)E_{12}(c_2)...E_{12}(c_m)E_{32}(d_1)...\widehat{E_{32}(d_i)}...E_{32}(d_n).1
\end{aligned}$$

$\blacksquare$

$$
\begin{aligned}
=&E_{21}(a_1)...E_{21}(a_m)E_{23}(b_1)...E_{23}(b_{n-1})E_{12}(c_1)E_{12}(c_2)...E_{12}(c_m)E_{23}(b_n)E_{32}(d_1)...E_{32}(d_n).1\\
&+\sum_{i=1}^{n}\sum_{j=1}^{m}E_{21}(a_1)...E_{21}(a_m)E_{23}(b_1)...E_{23}(b_{n-1}).\\
&E_{12}(c_1)...E_{12}(-c_jb_nd_i)...E_{12}(c_m)E_{32}(d_1)...\widehat{E_{32}(d_i)}...E_{32}(d_n).1\\
=&\sum_{i=1}^{n}\sum_{j>i}E_{21}(a_1)...E_{21}(a_m)E_{23}(b_1)...E_{23}(b_{n-1})E_{12}(c_1)E_{12}(c_2)...E_{12}(c_m)\\
&E_{32}(d_1)...\widehat{E_{32}(d_i)}...E_{32}(d_{j-1})E_{32}(-d_ib_nd_j-d_jb_nd_i)...E_{32}(d_n).1\\
&+\sum_{i=1}^{n}E_{21}(a_1)...E_{21}(a_m)E_{23}(b_1)...E_{23}(b_{n-1}).\\
&E_{12}(c_1)E_{12}(c_2)...E_{12}(c_m)E_{32}(d_1)...\widehat{E_{32}(d_i)}...E_{32}(d_n)(-\mu)\kappa(b_nd_i).1\\
&+\sum_{i=1}^{n}\sum_{j=1}^{m}E_{21}(a_1)...E_{21}(a_m)E_{23}(b_1)...E_{23}(b_{n-1}).\\
&E_{12}(c_1)...E_{12}(-c_jb_nd_i)...E_{12}(c_m)E_{32}(d_1)...\widehat{E_{32}(d_i)}...E_{32}(d_n).1.
\end{aligned}
$$

Using (3.4) is true for $n-1$, and expanding it we can get it is also true for n. $\square$

Lemma 3.6. *The hermitian form on different level is 0.*

Proof. Only need to prove those elements in the basis $\mathcal{B}$. Let

$$
u=E_{12}(a_1)...E_{12}(a_m)E_{32}(b_1)...E_{32}(b_n).1,\quad v=E_{12}(c_1)...E_{12}(c_k)E_{32}(d_1)...E_{32}(d_l).1,
$$

and $(m,n)\neq(k,l)$.

At first we prove $(u,v)=0$ with $m=0$:

If $k=0$, we can suppose $n>l$, then

$$
\begin{aligned}
(u,v)=&(E_{32}(b_1)...E_{32}(b_n).1,E_{32}(d_1)...E_{32}(d_l).1)\\
=&((-1)^lE_{23}(\bar{d_l})...E_{23}(\bar{d_1})E_{32}(b_1)...E_{32}(b_n).1,1)
\end{aligned}
$$

by Lemma 2.7, $lev(E_{23}(\bar{d_l})...E_{23}(\bar{d_1})E_{32}(b_1)...E_{32}(b_n).1)=(0,n-l)$ or $E_{23}(\bar{d_l})...E_{23}(\bar{d_1})E_{32}(b_1)...E_{32}(b_n).1=0$, then $(u,v)=0$.

For $k>0$,

$$
\begin{aligned}
(u,v)=&(E_{32}(b_1)...E_{32}(b_n).1,E_{12}(c_1)...E_{12}(c_k)E_{32}(d_1)...E_{32}(d_l).1)\\
=&(-E_{21}(\overline{c_1})E_{32}(b_1)...E_{32}(b_n).1,E_{12}(c_2)...E_{12}(c_k)E_{32}(d_1)...E_{32}(d_l).1)
\end{aligned}
$$

then from Lemma 2.7, $-E_{21}(\overline{c_1})E_{32}(b_1)...E_{32}(b_n).1 = 0$, then $(u,v) = 0$.

Without loss of generality, we can assume that $m \leq k$, then

$$\begin{aligned}&(u,v)\\ =&(E_{12}(a_1)...E_{12}(a_m)E_{32}(b_1)...E_{32}(b_n).1, E_{12}(c_1)...E_{12}(c_k)E_{32}(d_1)...E_{32}(d_l).1)\\ =&(E_{32}(b_1)...E_{32}(b_n).1, (-1)^m E_{21}(\overline{a_m})...E_{21}(\overline{a_1}).E_{12}(c_1)...E_{12}(c_k)E_{32}(d_1)...E_{32}(d_l).1).\end{aligned}$$

From Lemma 2.7,

$$lev(E_{21}(\overline{a_m})...E_{21}(\overline{a_1}).E_{12}(c_1)...E_{12}(c_k)E_{32}(d_1)...E_{32}(d_l).1) = (k-m, n)$$

or

$$E_{21}(\overline{a_m})...E_{21}(\overline{a_1}).E_{12}(c_1)...E_{12}(c_k)E_{32}(d_1)...E_{32}(d_l).1 = 0,$$

then back to the case $m = 0$, we get $(u,v) = 0$. $\square$

Similarly to [Proposition 4.11, GZ], and together with Lemma 3.3, we have

Proposition 3.7. *The hermitian form on the same element h in level (m,n) is a polynomial of μ, with the leading term is $c(-1)^{m+n}(-\mu)^{m+n} = c\mu^{m+n}$ with some constant $c > 0$.*

Now we can show the following theorem.

Theorem 3.8. *(π, V) is unitariazable if and only if $\mu > 0$.*

Proof. From [Theorem 4.12, GZ], the hermitian form in level $(0,n)$ and $(m,0)$ is positive definite if and only if needs $\mu > 0$.

Define

$$T_{a,b}(s^{m_1}t^{n_1}s^{m_2}t^{n_2}...s^{m_k}t^{n_k}) = s^{m_1+a}t^{n_1+b}s^{m_2+a}t^{n_2+b}...s^{m_k+a}t^{n_k+b}$$

$(a,b \in \mathbb{Z})$. Extend this operator to the linear operator $\widetilde{T_{a,b}}$ on V by

$$\begin{aligned}&\widetilde{T_{a,b}}(E_{12}(\alpha_1)E_{12}(\alpha_2)...E_{12}(\alpha_k)E_{32}(\beta_1)E_{32}(\beta_2)...E_{32}(\beta_l).1)\\ =&E_{12}(T_{a,b}\alpha_1)E_{12}(T_{a,b}\alpha_2)...E_{12}(T_{a,b}\alpha_k)E_{32}(T_{a,b}\beta_1)E_{32}(T_{a,b}\beta_2)...E_{32}(T_{a,b}\beta_l).1.\end{aligned}$$

Following Lemma 3.3, $\widetilde{T_{a,b}}$ preserves the hermitian form on V. Denote

$$\begin{aligned}L_{l,r}(M,N) = Span\{&E_{12}(s^{m_1}t^{n_1})...E_{12}(s^{m_l}t^{n_l})E_{32}(s^{j_1}t^{k_1})...E_{32}(s^{j_r}t^{k_r}).1\\ &|m_i, n_i \geqq 0, i = 1..l, j_\iota, k_\iota \geqq 0,\\ &\sum_{i=1}^{l} m_i + \sum_{\iota=1}^{r} j_\iota \leqq M, \sum_{i=1}^{r} n_i + \sum_{\iota=1}^{r} k_\iota \leqq N\}.\end{aligned}$$

Since the hermitian form on different level is 0, we will prove the unitarity by induction on the level.

For any $\mu > 0$, the form is definite in level $(0, n)$ ([Theorem 4.12,GZ]), and suppose it is definite in level (r, n), for those $r < m$, and it is not definite in level (m, n).

From Proposition 3.7, we know that the hermitian form restrict to this level should be positive definite for μ big enough. Assume it is not positive definite for some $\mu > 0$, then there exist M, N such that the form restrict on $L_{m,n}(M, N)$ is not positive definite. From Proposition 3.7, the form on $L_{l,r}(M, N)$ varies smoothly with μ. Then we can find a μ_0 at which the form is not positive definite, and for all $\mu > \mu_0$, it is positive definite. And we write $(.,.)_\mu$ to be the hermitian form at μ.

So the radical of the form is non-trivial at μ_0, i.e there exist a nonzero $\widetilde{h} \in L_{m,n}(M, N)$, such that for any $h \in L_{m,n}(M, N)$ we have

$$(\widetilde{h}, h)_{\mu_0} = 0.$$

Therefore for any arbitrary element $h_{m-1,n}$ in $L_{m-1,n}(M, N)$, and any $c \in \mathbb{C}$, we have

$$(E_{21}(c).\widetilde{h}, h_{m-1,n})_{\mu_0} = 0.$$

Since the form is positive definite in level $(m - 1, n)$, we have $E_{21}(c).\widetilde{h} = 0$, for any $c \in \mathbb{C}$. Replacing $\widetilde{h}$ by $\widetilde{T_{-a,-b}}(\widetilde{h})$ if necessary, we can write

$$\widetilde{h} = \sum_{i=1}^{m} a_i (E_{12}(1))^i x_i$$

where $x_i = \sum E_{12}(\alpha_{i+1})..E_{12}(\alpha_m)E_{32}(\beta_1)..E_{32}(\beta_n).1$ (here it is a finite sum), and α_i, β_j is the form $s^l t^k$ and l, k can not both be 0.

Let i_0 be the smallest one such that $a_{i_0} \neq 0$, then $i_0 \geq 1$.

Since

$$\begin{aligned}
&E_{21}(c)(E_{12}(1))^{i}E_{12}(\alpha_{i+1})..E_{12}(\alpha_m)E_{32}(\beta_1)..E_{32}(\beta_n).1\\
=&(E_{12}(1))^{i}E_{21}(c)E_{12}(\alpha_{i+1})..E_{12}(\alpha_m)E_{32}(\beta_1)..E_{32}(\beta_n).1\\
&+i.(E_{12}(1))^{i-1}(E_{22}(c)-E_{11}(c))E_{12}(\alpha_{i+1})..E_{12}(\alpha_m)E_{32}(\beta_1)..E_{32}(\beta_n).1\\
&+(-2c)\frac{i.(i-1)}{2}(E_{12}(1))^{i-1}E_{12}(\alpha_{i+1})..E_{12}(\alpha_m)E_{32}(\beta_1)..E_{32}(\beta_n).1\\
=&(E_{12}(1))^{i}E_{21}(c)E_{12}(\alpha_{i+1})..E_{12}(\alpha_m)E_{32}(\beta_1)..E_{32}(\beta_n).1\\
&+i.(E_{12}(1))^{i-1}((-2c)(m-i)-n)E_{12}(\alpha_{i+1})..E_{12}(\alpha_m)E_{32}(\beta_1)..E_{32}(\beta_n).1\\
&+i.(E_{12}(1))^{i-1}E_{12}(\alpha_{i+1})..E_{12}(\alpha_m)E_{32}(\beta_1)..E_{32}(\beta_n)(E_{22}(c)-E_{11}(c)).1\\
&+(-2c)\frac{i.(i-1)}{2}(E_{12}(1))^{i-1}E_{12}(\alpha_{i+1})..E_{12}(\alpha_m)E_{32}(\beta_1)..E_{32}(\beta_n).1\\
=&(E_{12}(1))^{i}E_{21}(c)E_{12}(\alpha_{i+1})..E_{12}(\alpha_m)E_{32}(\beta_1)..E_{32}(\beta_n).1\\
&+[ic(-\mu_0)+i((-2c)(m-i)-n)+(-2c)\frac{i.(i-1)}{2}].\\
&(E_{12}(1))^{i-1}E_{12}(\alpha_{i+1})..E_{12}(\alpha_m)E_{32}(\beta_1)..E_{32}(\beta_n).1,
\end{aligned}$$

we have

$$E_{21}(c)\widetilde{h}=\gamma a_{i_0}(E_{12}(1))^{i_0-1}x_{i_0}+R$$

where R contains those with power of $E_{12}(1)$ greater than i_0-1, and

$$\gamma=i_0c(-\mu_0)+i_0((-2c)(m-i_0)-n)+(-2c)\frac{i_0.(i_0-1)}{2}=ci_0(-\mu_0-(m-i_0)-(m-1)).$$

Since $m\geq i_0$, $i_0\geq 1$, and $\mu_0\geq 0$, $\gamma\neq 0$, contradict with $E_{21}(c)\widetilde{h}=0$.
So for any $\mu>0$, the hermitian form is positive definite. □

Acknowledgments

I am grateful to my supervisors Professors Nantel Bergeron and Yun Gao for their encouragement and support during the preparation of this paper, especially to Professor Yun Gao for drawing my attention to this subject.

REFERENCES

[AABGP] B. N. Allison, S. Azam, S. Berman, Y. Gao, A. Pianzola, *Extended affine Lie algebras and their root systems*, Memoir. Amer. Math. Soc. **126** (1997), Number 605.

[ABGP] B. N. Allison, S. Berman, Y. Gao, A. Pianzola, *A characterization of affine Kac-Moody Lie algebras*, Comm. Math. Phys. **185** (1997), 671–688.

[BEG] V. Baranovsky, S. Evens, V. Ginzburg, *Representations of quantum tori and double-affine Hecke algebras*, math.RT/0005024.

[BGK] S. Berman, Y. Gao, Y. Krylyuk, *Quantum tori and the structure of elliptic quasi-simple Lie algebras*, J. Funct. Anal. **135** (1996), 339–389.

[BGT] S. Berman, Y. Gao, S. Tan, *A Unified View of some Vertex Operator Constructions*, Israel J. Math. **134** (2003), 29–60.

[BS] S. Berman and J. Szmigielski, *Principal realization for extended affine Lie algebra of type sl_2 with coordinates in a simple quantum torus with two variables*, Cont.Math. **248** (1999), 39–67.

[BZ] Y. Billig, K. Zhao, *Vertex operator representations of quantum tori at roots of unity*, Commun. Contemp. Math. **6** (2004), 195–220.

[ER1] S. Eswara Rao, *A class of integrable modules for the core of EALA coordinatized by quantum tori*, J. Alg. **275** (2004), 59–74.

[ER2] S. Eswara Rao, *Unitary modules for EALAs co-ordinatized by a quantum torus*, Comm. Alg. **31** (2003), 2245–2256.

[EB] S. Eswara Rao, P. Batra, *A new class of representations of EALA coordinated by quantum tori in two variables. Dedicated to Robert V. Moody*, Canad. Math. Bull. **45** (2002), 672–685.

[EZ] S. Eswara Rao, K. Zhao, *Highest weight irreducible representations of rank 2 quantum tori*, Math. Res. Lett. **11** (2004), 615–628.

[EFK] P. Etingof and I. Frenkel, A. Kirillov, *Lectures on representation theory and Knizhnik-Zamolodchikov equations.*, Mathematical Surveys and Monographs, vol. 58, 1998.

[FF] B. Feigin and E. Frenkel, *Affine Kac-Moody algebras and semi-infinite flag manifolds*, Comm. Math. Phys. **128** (1990), 161–189.

[G1] Y. Gao, *Representations of extended affine Lie algebras coordinatized by certain quantum tori*, Compositio Mathematica **123** (2000), 1–25.

[G2] Y. Gao, *Vertex operators arising from the homogeneous realization for $\widehat{gl}_N$*, Comm. Math. Phys. **211** (2000), 745–777.

[G3] Y. Gao, *Fermionic and bosonic representations of the extended affine Lie algebra $\widetilde{\mathfrak{gl}_N(\mathbb{C}_q)}$*, Canada Math Bull. **45** (2002), 623–633.

[GZ] Y. Gao and Z. Zeng, *Hermitian representations of the extended affine Lie algebra $\widetilde{\mathfrak{gl}_2(\mathbb{C}_q)}$*, Adv. Math. (to appear).

[G-KK] M. Golenishcheva-Kutuzova and V. G. Kac, *Γ-conformal algebras*, J. Math. Phys. **39** (1998), 2290–2305.

[H-KT] R. Høegh-Krohn and B. Torresani, *Classification and construction of quasi-simple Lie algebras*, J. Funct. Anal. **89** (1990), 106–136.

[JK1] H. P. Jakobsen and V. G. Kac, *A new class of unitarizable highest weight representations of infinite-dimensional Lie algebras.*, Lecture Notes in Physics **226** (1985), 1–20.

[JK2] H. P. Jakobsen and V. G. Kac, *A new class of unitarizable highest weight representations of infinite-dimensional Lie algebras. II*, J. Funct. Anal. **82** (1989), 69–90.

[L] M. Lau, *Bosonic and fermionic representations of Lie algebra central extensions*, Adv. Math. **194** (2005), 225–245.

[LT1] W. Lin, S. Tan, *Representations of the Lie algebra of derivations for quantum torus*, J. Alg. **275** (2004), 250–274.

[LT2] W. Lin, S. Tan, *Harish-Chandra modules for the q-analog Virasoro-like algebra*, J. Alg. (to appear).

[M] Y. I. Manin, *Topics in noncommutative geometry*, Princeton University Press, 1991.

[Mi] K. Miki, *Integrable irreducible highest weight modules for* $sl_2(C_p[x^{\pm 1}, y^{\pm 1}])$, Osaka J. Math. **41** (2004), 295–326.

[S] K. Saito, *Extended affine root systems 1 (Coxeter transformations)*, Publ. RIMS., Kyoto Univ. **21** (1985), 75–179.

[SZ] Y. Su, L. Zhu, *Representations of the Derivation Algebra of the Localization of the Quantum Plane at* $q = -1$, Comm. Alg. (to appear).

[VV] M. Varagnolo and E. Vasserot, *Double-loop algebras and the Fock space*, Invent. Math. **133** (1998), 133–159.

[W1] M. Wakimoto, *Extended affine Lie algebras and a certain series of Hermitian representations*, Preprint (1985).

[W2] M. Wakimoto, *Representations of the affine Lie algebra* $A_1^{(1)}$, Comm. Math. Phys. **104** (1986), 605–609.

[ZZ] H. Zhang, K. Zhao, *Representations of the Virasoro-like algebra and its q-analog*, Comm. Algebra **24** (1996), 4361–4372.

DEPARTMENT OF MATHEMATICS AND STATISTICS, YORK UNIVERSITY, TORONTO,
CANADA M3J 1P3
E-mail address: `ziting@mathstat.yorku.ca`